

SMITHSONIAN
LIBRARIES



SHOUT
SHARE **IT**





Oct 15 26 3096 Smith 76

Education Department Bulletin

Published fortnightly by the University of the State of New York

Entered as second-class matter June 24, 1908, at the Post Office at Albany, N. Y., under the act of July 16, 1894

No. 470

ALBANY, N. Y.

MAY 1, 1910

New York State Museum


JOHN M. CLARKE, Director

CHARLES H. PECK, State Botanist

Museum Bulletin 139

REPORT OF THE STATE BOTANIST 1909

	PAGE		PAGE
Introduction.....	5	List of edible, poisonous and	
Plants added to the herbarium...	8	unwholesome mushrooms	
Contributors and their contribu-		hitherto figured and described	
tions	10	by C. H. Peck.....	78
Species not before reported.....	19	List of genera whose New York	
Remarks and observations.....	33	species (chiefly) have been col-	
Edible fungi.....	37	lated with descriptions in the	
New species of extralimital fungi.	42	State Botanist's reports cited..	87
New York species of <i>Inocybe</i> ...	48	Explanation of plates	91
New York species of <i>Hebeloma</i> ..	67	Index.....	III



ALBANY

UNIVERSITY OF THE STATE OF NEW YORK

1910

STATE OF NEW YORK
EDUCATION DEPARTMENT

Regents of the University
With years when terms expire

1913	WHITELAW REID M.A. LL.D. D.C.L.	<i>Chancellor</i>	New York
1917	ST CLAIR MCKELWAY M.A. LL. D.	<i>Vice Chancellor</i>	Brooklyn
1919	DANIEL BEACH Ph.D. LL.D.	- - - - -	Watkins
1914	PLINY T. SEXTON LL.B. LL.D.	- - - - -	Palmyra
1912	T. GUILFORD SMITH M.A. C.E. LL.D.	- - - - -	Buffalo
1918	WILLIAM NOTTINGHAM M.A. Ph.D. LL.D.	- - - - -	Syracuse
1922	CHESTER S. LORD M.A. LL.D.	- - - - -	New York
1915	ALBERT VANDER VEER M.D. M.A. Ph.D. LL.D.	- - - - -	Albany
1911	EDWARD LAUTERBACH M.A. LL.D.	- - - - -	New York
1920	EUGENE A. PHILBIN LL.B. LL.D.	- - - - -	New York
1916	LUCIAN L. SHEDDEN LL.B. LL.D.	- - - - -	Plattsburg
1921	FRANCIS M. CARPENTER - - - - -	- - - - -	Mount Kisco

Commissioner of Education

ANDREW S. DRAPER LL.B. LL.D.

Assistant Commissioners

AUGUSTUS S. DOWNING M.A. Pd.D. LL.D. *First Assistant*

FRANK ROLLINS Ph.D. *Second Assistant*

THOMAS E. FINEGAN M.A. Pd.D. *Third Assistant*

Director of State Library

JAMES I. WYER, JR, M.L.S.

Director of Science and State Museum

JOHN M. CLARKE Ph.D. Sc.D. LL.D.

Chiefs of Divisions

Administration, HARLAN H. HORNER B.A.

Attendance, JAMES D. SULLIVAN

Educational Extension, WILLIAM R. EASTMAN M.A. M.L.S.

Examinations, CHARLES F. WHELOCK B.S. LL.D.

Inspections, FRANK H. WOOD M.A.

Law, FRANK B. GILBERT B.A.

School Libraries, CHARLES E. FITCH L.H.D.

Statistics, HIRAM C. CASE

Trades Schools, ARTHUR D. DEAN B.S.

Visual Instruction, ALFRED W. ABRAMS Ph.B.

207
New York State Education Department

Science Division, February 23, 1910

Hon. Andrew S. Draper LL.D.

Commissioner of Education

SIR: I have the honor to communicate herewith for publication as a bulletin of the State Museum, the report of the State Botanist for the fiscal year ending September 30, 1909.


Very respectfully

JOHN M. CLARKE

Director

State of New York
Education Department
COMMISSIONER'S ROOM

Approved for publication this 24th day of February 1910

A large, stylized handwritten signature in dark ink, appearing to read 'A. V. Conner'. The signature is written over a horizontal line and has a long, sweeping flourish extending downwards and to the right.

Commissioner of Education

Education Department Bulletin

Published fortnightly by the University of the State of New York

Entered as second-class matter June 24, 1908, at the Post Office at Albany, N. Y., under the act of July 16, 1894

No. 470

ALBANY, N. Y.

#

MAY 1, 1910

New York State Museum

JOHN M. CLARKE, Director
CHARLES H. PECK, State Botanist

Museum Bulletin 139

REPORT OF THE STATE BOTANIST 1909

Dr John M. Clarke, Director of State Museum:

The following report of work done in the botanical department of the State Museum for the year 1909 is respectfully submitted.

Since the date of my last report specimens of plants for the State herbarium have been collected in the counties of Albany, Columbia, Jefferson, Lewis, Livingston, Rensselaer, Steuben, St Lawrence, Warren and Wyoming. Specimens have also been added to the herbarium that were received from correspondents and others. These were collected in the counties of Albany, Cayuga, Dutchess, Essex, Franklin, Herkimer, Monroe, New York, Oneida, Onondaga, Ontario, Orleans, Oswego, Queens, Rensselaer, Schoharie, Suffolk, Tompkins, Ulster, Warren and Washington.

The number of species of which specimens have been added to the herbarium is 255 of which 56 species were not before represented in it. Of these, 11 are considered new or hitherto undescribed species. All except one are fungi. The specimens of the 199 species not new to the herbarium serve to give a better or more complete representation of their respective species than was given before. A list of the names of all the added species is given under the title "Plants added to the herbarium."

The number of those reported as contributors to the herbarium is 66. Some of these have sent specimens for identification merely, but when the specimens were collected in this State and were received in good condition, if the species was previously unrepresented in the herbarium or if for any other reason they were deemed worthy of preservation, they have been preserved and credited to the sender as a contribution to the herbarium.

Some of our best and most interesting additions to the herbarium have been made in this way. The names of contributors of such specimens and of extralimital specimens with their respective contributions are given under the title "Contributors and their contributions."

The number of species added to our New York flora is 77. Several of these have been reported before as varieties of other species or confused with other species, but having been recently admitted in Gray's New Manual as distinct species it has been thought best to record them as such with their known New York localities. The names of these and other added species together with their localities, descriptions of new species, and other matters of interest will be found under the title "Species not before reported." Under the heading "Remarks and observations" any facts of interest concerning the species mentioned are recorded. This record may include new varieties of plants or notable variations, new localities for rare plants, and remarks concerning the diseases of plants or their economic properties.

The work of testing our wild mushrooms for their edible qualities as opportunity was given has been continued. Five species have been personally tested and approved as edible. These, added to the species and varieties previously known, make the number of New York species and varieties now known to be edible 200. Plain and simple descriptions of the newly added species are given under the title "Edible fungi." Colored figures of these species may be found on plates 117-20. Among the extralimital contributed specimens 10 apparently new species are represented. Descriptions of these species are given in a part of the report marked "New species of extralimital fungi." Colored figures of six of these species may be found on plates W, X, Y and Z.

Revised descriptions of our New York species of the genera *Inocybe* and *Hebeloma* have been prepared, with keys to the sections or subgenera and to the species. It is believed that these simple localized monographs will be helpful to those studying or desiring to study these interesting subjects of the vegetable kingdom. These chapters are respectively entitled "New York species of *Inocybe*" and "New York species of *Hebeloma*."

The climatic character of the season has been to a large extent a repetition of that of 1908. A cold late spring, an unusually dry summer and prevailing cool weather were its characteristic

features. These conditions were decidedly unfavorable to wild mushroom growth. Scarcely any could be found except the few that naturally inhabit swamps and low wet ground in woods. In the latter part of the season gentle rains moistened the surface of the ground sufficiently to insure the development of good agricultural crops and a fair seasonable crop of some mushrooms. But the effect upon the common mushroom, *Agaricus campester* and its variety, the garden mushroom, *Agaricus campester hortensis*, is worthy of special notice. In the vicinity of Albany a gentle and prolonged rain, the latter part of August, moistened the surface of the ground quite effectually. In a few days the common mushroom appeared in unusual abundance, though it was a little earlier in the season than it usually appears. The mushrooms were so plentiful that at least one fruit dealer offered them for sale in quart baskets at his fruit stand. A few weeks later light showers were followed by a copious crop of the "garden mushroom," a form differing from the common mushroom in having its cap adorned with brownish fibrils which form small spotlike scales on it and give it a darker color than that of the white form of the common mushroom. This crop continued to develop freely for several days and grew in some instances in pastures of light sandy soil where mushrooms are not usually expected to grow. The same abundant appearance of the edible mushroom was reported to have followed the light autumnal showers in other localities in the State. The lesson it teaches is that for mushroom production gentle showers are better than torrents of rain.

The number of those who have sent or brought specimens of plants to the office of the botanist for identification is 152. The number of identifications made is 1717.

Mr S. H. Burnham, my assistant, in addition to his other duties, has prepared a list of the names of the edible, poisonous and unwholesome species of mushrooms hitherto figured and described in the publications of the museum, together with the citations of the time and place of publication of each. He has also prepared a list of the genera of fungi of which the New York species (chiefly) have been described as far as known in previous reports. The time and place of these limited monographic publications are cited. Both these lists may be found at the end of this report.

CHARLES H. PECK

State Botanist

Albany, December 24, 1909

PLANTS ADDED TO THE HERBARIUM

New to the herbarium

- | | |
|--|---|
| Ascochyta solani-nigri <i>Diedicke</i> . | Microcera coccophila <i>Desm.</i> |
| Belonidium glyceriae <i>Pk.</i> | Midotis irregularis (<i>Schw.</i>) |
| Biatora cupreo-rosella (<i>Nyl.</i>) <i>Tuckerm.</i> | Monolepis nuttalliana (<i>R. & S.</i>) |
| Bidens tenuisecta <i>Gray</i> | Morchella crispa <i>Karst.</i> |
| Boletus viridarius <i>Frost</i> | M. rimosipes <i>DC.</i> |
| Carduus crispus <i>L.</i> | Nardia crenulata (<i>Sw.</i>) <i>Lindb.</i> |
| Chaenactis stevioides <i>H. & A.</i> | N. hyalina (<i>Lyell</i>) <i>Carr.</i> |
| Ciboria luteo-virescens <i>R. & D.</i> | Peridermium strobili <i>Kleb.</i> |
| Clitocybe candida <i>Bres.</i> | Pezizella lanc.-paraphysata <i>Rehm</i> |
| Cortinarius subsalmoneus <i>Kauf. Ms.</i> | Phaeopezia fuscocarpa (<i>E. & H.</i>) |
| Crataegus brevipes <i>Pk.</i> | Pholiota aurivella <i>Batsch</i> |
| C. efferata <i>S.</i> | Phomopsis stewartii <i>Pk.</i> |
| C. letchworthiana <i>S.</i> | Picris echinoides <i>L.</i> |
| Diplocladium penicilloides <i>Sacc.</i> | Polyporus giganteus (<i>Pers.</i>) <i>Fr.</i> |
| Diplodia cercidis <i>E. & E.</i> | Psilocybe nigrella <i>Pk.</i> |
| D. hamamelidis <i>Fairm.</i> | Puccinia epiphylla (<i>L.</i>) <i>Wettst.</i> |
| D. tamariscina <i>Sacc.</i> | Ribes trist. albinervium (<i>Mx.</i>) |
| Dothiorella divergens <i>Pk.</i> | Rubia tinctorum <i>L.</i> |
| Epipactis tessellata (<i>Lodd.</i>) <i>Eaton</i> | Rumex pallidus <i>Bigel.</i> |
| Fenestella amorphia <i>E. & E.</i> | Schwalbea americana <i>L.</i> |
| Hypholoma boughtoni <i>Pk.</i> | Septoria sedicola <i>Pk.</i> |
| H. rigidipes <i>Pk.</i> | Solidago aspera <i>Ait.</i> |
| Leontodon nudicaulis (<i>L.</i>) <i>Banks</i> | Sparganium diversifolium <i>Graeb.</i> |
| Ligusticum scoticum <i>L.</i> | Stachys sieboldii <i>Miq.</i> |
| Lophiotrema hysterioides <i>E. & E.</i> | Stephanoma strigosum <i>Wallr.</i> |
| L. littorale <i>Speg.</i> | Trametes merisma <i>Pk.</i> |
| Marasmius alienus <i>Pk.</i> | Verticillium rexianum <i>Sacc.</i> |
| Melanopsamma confertissima (<i>Plow.</i>) | Volvaria volvacea (<i>Bull.</i>) <i>Fr.</i> |

Not new to the herbarium

- | | |
|---|---|
| Agaricus campester <i>L.</i> | Aristida gracilis <i>Ell.</i> |
| A. silvicola <i>Vitt.</i> | Armillaria mellea <i>Vahl</i> |
| Agropyrum tenerum <i>Vasey</i> | Aster ericoides <i>L.</i> |
| Alnus crispa (<i>Ait.</i>) <i>Pursh</i> | A. puniceus <i>L.</i> |
| Amanita frostiana <i>Pk.</i> | Atriplex pat. littoralis (<i>L.</i>) |
| A. phalloides <i>Fr.</i> | Barbarea vulgaris <i>R. Br.</i> |
| Anagallis arvensis <i>L.</i> | Bidens beckii <i>Torr.</i> |
| Angelica atropurpurea <i>L.</i> | Boehmeria cylindrica (<i>L.</i>) <i>Sw.</i> |
| Antennaria brainerdi <i>Fern.</i> | Bromus altissimus <i>Pursh</i> |
| Anthemis cotula <i>L.</i> | Caldesiella ferruginosa (<i>Fr.</i>) |
| Arcyria cinerea (<i>Bull.</i>) <i>Pers.</i> | Calvatia elata (<i>Mass.</i>) <i>Morg.</i> |
| A. punicea <i>Pers.</i> | C. gigantea (<i>Batsch</i>) |
| Arenaria peploides <i>L.</i> | Cantharellus infundibuliformis (<i>Scop.</i>) |
| Arisaema dracontium (<i>L.</i>) <i>Schott</i> | Cardamine bulbosa (<i>Schreb.</i>) |

- Cardamine douglassii* (Torr.)
Carduus spinosissimus (Walt.)
Carex bebbii Olney
C. crawfordii Fern.
Centaurea solstitialis L.
Cerastium viscosum L.
Cladosporium typhae Schw.
Collybia myriadophylla Pk.
C. platyphylla Fr.
C. radicata (Relh.) Fr.
Convallaria majalis L.
Coprinus atramentarius (Bull.)
C. micaceus (Bull.)
Cornus amomum Mill.
Corallorrhiza trifida Chat.
Crataegus acclivis S.
C. anomala S.
C. beata S.
C. eatoniana S.
C. ellwangeriana S.
C. grayana Eggl.
C. halliana S.
C. holmesiana Ashe
C. ignea S.
C. maineana S.
C. menandiana S.
C. ovatifolia S.
C. persimilis S.
C. polita S.
C. punctata Jacq.
C. repulsans S.
C. rotundifolia Moench
C. succulenta Lk.
C. tenuiloba S.
Crepidotus applanatus (Pers.)
C. malachius B. & C.
Cuscuta arvensis Beyrich.
C. cephalanthi Engelm.
Cypripedium acaule Ait.
Cystopus candidus (Pers.) Lev.
Daedalea unicolor (Bull.) Fr.
Erysiphe cichoracearum DC.
Erythronium albidum Nutt.
Eupatorium purpureum L.
Exidia gland. levior Sacc.
Exoascus confusus Atk.
E. pruni Fckl.
E. unilateralis Pk.
Fagopyrum tataricum (L.) Gaertn.
Favolus europaeus Fr.
Fimbristylis castanea (Mx.) Vahl
Flammula pulchrifolia Pk.
Fuligo septica (Lk.) Gmel.
Fusicladium destruens Pk.
Galera lat. albicolor Pk.
Galium erectum Huds.
Geranium pusillum Burm. f.
Geum flavum (Port.) Bickn.
Gloeoporus conchoides Mont.
Hedeoma hispida Pursh
Helenium autumnale L.
Hieracium florentinum All.
H. gronovii L.
H. scabrum Mx.
Hybanthus concolor (Forst.)
Hypericum canadense Mx.
Hypochaeris radicata L.
Ilex vert. tenuifolia (Torr.) Wats.
Iris prismatica Pursh
Irpex obliquus (Schrud.) Fr.
Juncus brachycephalus (Engelm.)
J. brevicaudatus (Engelm.)
J. secundus Beauv.
Juniperus com. depressa Pursh
J. horizontalis Moench
Lactarius aquifluus Pk.
L. glyciosmus Fr.
Lactuca scar. integrata G. & G.
Laportea canadensis (L.) Gaud.
Lappula virginiana (L.) Greene
Leonurus cardiaca L.
Listera australis Lindl.
Lycoperdon gemmatum Batsch
Marasmius acerinus Pk.
M. glabellus Pk.
M. oreades Fr.
Monilia crataegi Diedicke
Morus rubra L.
Mycena pelianthina Fr.
M. pseudopoda (Pers.)
M. pseudopura Cke.
M. sanguinolenta A. & S.
Myosotis virginica (L.) B. S. P.
Naias gracillima (A. Br.) Magn.
Oenothera linearis Mx.
Omphalia rugosodisca Pk.
Onopordon acanthium L.
Panicum implicatum Scribn.
P. oricola H. & C.
P. spretum Schultes

- Paxillus involutus (Batsch) Fr.*
Peridermium consimile A. & K.
Phlox divaricata L.
Pholiota angustipes Pk.
P. vermiflua Pk.
Pilea pumila (L.) Gray
Plantago decipiens Braineoud
Pleurotus ulmarius (Bull.) Sow.
Pluteus admirabilis Pk.
P. cervinus (Schaeff.) Fr.
P. granularis Pk.
P. nanus (Pers.) Fr.
Polygonum avic. littorale (Lk.)
Polyporus elegans Fr.
P. sulphureus (Bull.) Fr.
Potamogeton richardsoni (Benn.)
Prunus pumila L.
Puccinia coronata Cda.
P. rubigo-vera (DC.) Wint.
P. veratri Niessl.
Pyrus coronaria L.
P. melanocarpa (Mx.) Willd.
Quercus macrocarpa Mx.
Radicula pal. hispida (Desv.)
Ranunculus delphinifolius Torr.
R. reptans L.
Roestelia aurantiaca Pk.
Rubus andrewsianus Blanch.
R. permixtus Blanch.
R. recurvans Blanch.
Russula brevipes Pk.
R. lepida Fr.
R. mariae Pk.
Sanicula canadensis L.
Sedum ternatum Mx.
Silybium marianum (L.) Gaertn.
Sisymbrium altissimum L.
S. sophia L.
Solanum dulcamara L.
S. nigrum L.
Solidago neglecta T. & G.
S. squarrosa Muhl.
Sparganium americanum Nutt.
S. angustifolium Mx.
Spiraea latifolia Borkh.
Stachys arenicola Britton
Suaeda maritima (L.) Dumort.
Thalictrum confine Fern.
T. dasycarpum F. & L.
T. revolutum DC.
Thelephora terrestris Ehrh.
Trametes suaveolens (L.) Fr.
Tricholoma album (Schaeff.) Fr.
Trichothecium roseum (Pers.) Lk.
Trillium grand. variegatum Pk.
Tripsacum dactyloides L.
Urtica lyellii Wats.
Ustilago longissima (Sow.) Tul.
Ustulina vulgaris Tode
Vaccinium pennsylvanicum Lam.
Veronica humifusa Dicks.
V. tournefortii Gmel.
Verticillium lactarii Pk.
Vicia angustifolia (L.) Reich.
Viola rafinesquii Greene
V. renifolia Gray
V. sororia Willd.
V. triloba Schw.
Vitis vulpina L.
Zizania palustris L.
Zizia aurea (L.) Koch.

CONTRIBUTORS AND THEIR CONTRIBUTIONS

Miss L. C. Allen, Newtonville, Mass.

Bovistella ohiensis E. & M.

Miss H. C. Anderson, Lambertville, N. J.

Morchella gigas (Batsch) Fr.

Miss F. Beckwith, Rochester

Bidens tenuisecta Gray

Chaenactis stevioides H. & A.

Erodium cicutarium (L.) L'Her.

Geranium pusillum Burm. f.

Monolepis nuttalliana (R. & S.)

Sisymbrium sophia L.

Viola sororia Willd.

Mrs E. B. Blackford, Boston, Mass.

- | | |
|----------------------------------|---------------------------------|
| Cortinarius acutoides <i>Pk.</i> | Lactarius hyssiginus <i>Fr.</i> |
| C. lutescens <i>Pk.</i> | Russula blackfordae <i>Pk.</i> |
| | Russula serissima <i>Pk.</i> |

Mrs H. C. Davis, Falmouth, Me.

- | | |
|---------------------------------|--|
| Bovista pila <i>B. & C.</i> | Mutinus caninus (<i>Huds.</i>) <i>Fr.</i> |
| Crucibulum vulgare <i>Tul.</i> | Rhizina inflata (<i>Schaeff.</i>) <i>Quel.</i> |
- A set of colored drawings representing about 150 species of fleshy fungi from Maine

Mrs E. P. Gardner, Canandaigua

- Trillium grandiflorum variegatum *Pk.*

Mrs L. L. Goodrich, Syracuse

- | | |
|---|---------------------------------|
| Arisaema dracontium (<i>L.</i>) <i>Schott</i> | Sisymbrium altissimum <i>L.</i> |
| Veronica tournefortii <i>C. C. Gmeln.</i> | |

Mrs C. W. Harris, Washington, D. C.

- | | |
|--|---|
| Baeomyces roseus <i>Pers.</i> | Peltigera canina (<i>L.</i>) <i>Hoffm.</i> |
| Cetraria oakesiana <i>Tuckm.</i> | P. polydactyla (<i>Neck.</i>) |
| Cladonia caespiticia (<i>Pers.</i>) <i>Fl.</i> | Physcia aquila detonsa <i>Tuckm.</i> |
| C. cristatella <i>Tuckm.</i> | P. caesia (<i>Hoffm.</i>) <i>Nyl.</i> |
| C. mitrula <i>Tuckm.</i> | P. obscura (<i>Ehrh.</i>) <i>Nyl.</i> |
| C. papillaria (<i>Ehrh.</i>) <i>Hoffm.</i> | P. obsc. endochrysea <i>Nyl.</i> |
| C. pyxidata (<i>L.</i>) <i>Fr.</i> | P. stellaris (<i>L.</i>) <i>Tuckm.</i> |
| C. rangiferina (<i>L.</i>) <i>Hoffm.</i> | P. stell. aipolia <i>Nyl.</i> |
| C. verticillata <i>Fr.</i> | Pyxine soorediata <i>Fr.</i> |
| Parmelia borrieri rudecta <i>Tuckm.</i> | Ramalina calic. fastigiata <i>Fr.</i> |
| P. caperata (<i>L.</i>) <i>Ach.</i> | Sticta amplissima (<i>Scop.</i>) <i>Mass.</i> |
| P. conspersa (<i>Ehrh.</i>) <i>Ach.</i> | S. pulmonaria (<i>L.</i>) <i>Ach.</i> |
| P. perlata (<i>L.</i>) <i>Ach.</i> | Umbilicaria dillenii <i>Tuckm.</i> |
| P. physodes (<i>L.</i>) <i>Ach.</i> | U. muhlenbergii (<i>Ach.</i>) |
| P. saxatilis (<i>L.</i>) <i>Fr.</i> | Umbilicaria pustulata papulosa |
| Peltigera aphthosa (<i>L.</i>) <i>Hoffm.</i> | <i>Tuckm.</i> |

Miss A. Hibbard, West Roxbury, Mass.

- | | |
|--|----------------------------------|
| Boletinus glandulosus <i>Pk.</i> | Gomphidius nigricans <i>Pk.</i> |
| Boletus miniato-olivaceus <i>Frost</i> | Stropharia depilata <i>Pers.</i> |
| | Tricholoma acre <i>Pk.</i> |

Miss D. Hone, Minneapolis, Minn.

- | | |
|-----------------------------------|--------------------------------|
| Polyporus isidioides <i>Berk.</i> | Polyporus obtusus <i>Berk.</i> |
|-----------------------------------|--------------------------------|

Miss A. Lorenz, Hartford, Conn.

- | | |
|---|--|
| Marsupella robusta (<i>DeNot.</i>) <i>Evans</i> | Nardia crenulata (<i>Sm.</i>) <i>Lindb.</i> |
| M. sullivantii (<i>DeNot.</i>) | N. hyalina (<i>Lyell</i>) <i>Carr.</i> |

Miss H. L. Palliser, Poughkeepsie

- Boletus viridarius *Frost*

Dr C. E. Putnam, St Paul, Minn.

Secotium acuminatum Mont.

Miss M. L. Sutliff, Sacramento, Cal.

Rhizopogon rubescens Tul.

Mrs M. E. Williams, Wernersville, Pa.

Leskea gracilescens Hedw.

J. C. Arthur, Lafayette, Ind.

Puccinia grindeliae Pk.

A. D. Baker, Auburn

Centaurea solstitialis L.

C. F. Baker, Claremont, Cal.

<i>Agaricus bivelatus</i> Pk.	<i>Hypholoma campanulata</i> Pk.
A. <i>solidipes</i> Pk.	H. <i>cutifracta</i> Pk.
A. <i>subnitens</i> Pk.	<i>Inocybe bakeri</i> Pk.
<i>Amanita bivelvata</i> Pk.	I. <i>bulbosa</i> Pk.
A. <i>calyptratoides</i> Pk.	<i>Lactarius rufulus</i> Pk.
A. <i>ocreata</i> Pk.	L. <i>theiogalus</i> (Bull.)
A. <i>virosa</i> Fr.	<i>Leptonia edulis</i> Pk.
<i>Amanitopsis velosa</i> Pk.	<i>Mycena atroalboides</i> Pk.
<i>Armillaria subannulata</i> Pk.	M. <i>elegantula</i> Pk.
<i>Boletus tomentipes</i> Earle	M. <i>haematopoda</i> (Pers.) Fr.
<i>Clitocybe microspora</i> Pk.	<i>Naucoria platysperma</i> Pk.
C. <i>sphaerospora</i> Pk.	N. <i>vinicolor</i> Pk.
<i>Collybia albogrisea</i> Pk.	<i>Pluteolus luteus</i> Pk.
<i>Coprinus calyptratus</i> Pk.	<i>Psathyrella graciloides</i> Pk.
<i>Cortinarius multififormis</i> Fr.	<i>Psilocybe castanella</i> Pk.
<i>Hebeloma foedatum</i> Pk.	<i>Russula semicrema</i> Fr.
H. <i>ischnostylum</i> Cke.	<i>Tricholoma equestre</i> (L.) Fr.
	<i>Tubaria furfuracea</i> (Pers.) Fr.

H. J. Banker, Greencastle, Ind.

Onygena equina Pers. *Polyporus sulphureus* (Bull.) Fr.

Xylaria pedunculata (Dicks.) Fr.

H. W. Barratt, Poughkeepsie

Coprinus atramentarius (Bull.) Fr.

E. Bartholomew, Stockton, Kan.

- Barlaea subaurantia* B. & R.
Bjerkandera adusta (Willd.) Karst.
Botrytis uredinicola Pk.
Bubakia crotonis (Cke.) Arth.
Ceratophorum uncinatum (Clint.)
Cercospora biformis Pk.
C. *brunnea* Pk.
C. *flagellaris* E. & M.
C. *fuscovirens* Sacc.
C. *mississippiensis* T. & E.
C. *rhoina* C. & E.
C. *rubi* Sacc.
C. *simulata* E. & E.
C. *sordida* Sacc.
C. *vignae* E. & E.
Clavaria aurea Schaeff.
Coleosporium elephantopodis (Schw.)
C. *ipomoeae* (Schw.) Burr.
C. *laciniariae* Arth.
C. *solidaginis* (Schw.)
C. *vernoniae* B. & C.
Coriolus prolificans (Fr.) Murr.
C. *versicolor* (L.) Quel.
Corticium roseolum Mass.
Cylindrosporium padi Karst.
Daedalea aesculi (Schw.) Murr.
Darlucula filum (Biv.) Cast.
Diatrype stigma (Hoffm.) Fr.
Fusarium bartholomaei Pk.
F. *juglandinum* Pk.
Ganoderma curtisii (Berk.) Murr.
Gyroceras divergens Pk.
Hapalopilus gilvus (Schw.) Murr.
Helminthosporium hamatellum Pk.
Herpotrichia rhodospiloides Pk.
Hirneola auricula-judae (L.)
Hydnoporia fuscescens (Schw.) Murr.
Hypoxyton multifforme Fr.
Irpiciporus lacteus (Fr.) Murr.
Kuehneola albida (Kuehn.) Magn.
Lenzites betulina (L.) Fr.
Lycoperdon atropurpureum Vitt.
L. *pulcherrimum* B. & C.
Melampsora bigelowii Thuem.
M. *medusae* Thuem.
Microsphaera alni (Wallr.) Salm.
Nectria cinnabarina (Tode) Fr.
Nummularia repanda (Fr.) Nits.
Ozonium auricomum Link
Peniophora quercina (Fr.) Cke.
Phlebia radiata Fr.
Phyllosticta smilacis E. & M.
Piggotia fraxini B. & C.
Pileolaria toxicodendri (B. & R.)
Puccinia helianthi Schw.
P. *lateripes* B. & R.
P. *lobeliae* Ger.
P. *menthae americana* Pk.
P. *Muhlenbergiae* A. & H.
P. *polygoni-amphibii* Pers.
P. *smilacis* Schw.
P. *xanthii* Schw.
Pucciniastrum agrimoniae (Schw.)
P. *hydrangeae* (B. & C.)
P. *myrtilli* (Schw.)
Rhysotheca halstedii (Farl.)
Schizophyllum commune Fr.
Scleroderma tenerum B. & C.
Septoria musiva Pk.
S. *populi* Desm.
S. *rubi* West.
S. *scrophulariae* Pk.
Sorosporium ellisii Wint.
Sphaerella fraxinicola (Schw.)
Sphaeria potentillae Schw.
Stereum acerinum nivolum Berk.
S. *complicatum* Fr.
S. *curtisii* Berk.
S. *spadiceum* Fr.
S. *versicolor* (Sw.) Fr.
Stigmina platani (Fckl.) Sacc.
Thelephora rosella Pk.
Tranzschelia punctata (Pers.)
Uncinula parvula C. & P.
Uromyces andropogonis Tracy
U. *appendiculatus* (Pers.)
U. *aristidae* E. & E.
U. *euphorbiae* C. & P.
U. *hedysari-paniculata* (Schw.)
U. *lespedezae* (Schw.) Pk.
U. *spermacocis* (Schw.) Curt.
Valsaria nigrospora (Pk.) B. & V.

J. B. Bartlett, Albany

Sterigmatocystis ochracea (Wilh.) VanTigh.

F. S. Boughton, Pittsford*Hypholoma boughtoni* *Pk.**Volvaria volvacea* (*Bull.*) *Fr.***S. H. Burnham, Sandy Hill***Alnus crispa* (*Ait.*) *Pursh**Hypomyces lactifluorum* (*Schw.*)*Anthemis cotula* *L.**H. torminosus* (*Mont.*) *Tul.**Aster divaricatus* *L.**Julella monosperma* (*Pk.*) *Sacc.**A. , macrop. velutinus* *Bu.**Lactarius cinereus* *Pk.**Carduus crispus* *L.**L. subdulcis* (*Bull.*) *Fr.**Ceratiomyxa fruticulosa* (*Muell.*)*Lecanora rubina* (*Vill.*) *Ach.**Cladosporium herbarum* (*Pers.*) *Fr.**Massaria vomitoria* *B. & C.**Clitocybe candida* *Bres.**Peridermium conorum-piceae* (*Rees*)*C. trullisata* *Ellis**Peronospora parasitica* (*Pers.*)*Collybia platyphylla* *Fr.**Piggotia astroidea* *B. & Br.**Coprinus insignis* *Pk.**Polyporus chioneus* *Fr.**Corticium cremicolor* *B. & C.**Psilocybe uda* (*Pers.*) *Fr.**C. lacteum* *Fr.**Pucciniastrum potentillae* *Kom.**Cortinarius rimosus* *Pk.**Rubia tinctorum* *L.**C. subsalmoneus* *Kauff. Ms.**Rubus permixtus* *Blanch.**C. validipes* *Pk.**Russula aeruginea* *Fr.**Diplocladium penicilloides* *Sacc.**R. decolorans* *Fr.**Eutypella cerviculata* (*Fr.*) *Sacc.**Sanicula canadensis* *L.**Flammula pulchrifolia* *Pk.**Sparganium diversifolium* *Griseb.**F. spumosa* *Fr.**Stachys arenicola* *Britt.**Fomes pinicola* (*Sw.*) *Fr.**S. sieboldii* *Miq.**Geum flavum* (*Port.*) *Bickn.**Trametes sepium* *Berk.**Gloeosporium irregulare* *Pk.**T. suaveolens* (*L.*) *Fr.**Helvella palustris* *Pk.**Tricholoma mutans* *Pk.**Hydnum laciniatum* *Leers**Vaccinium pennsylvanicum* *Lam.**Hypocrea aurantiaca* *Pk.**Zygodesmus fuscus* *Corda***I. O. Cross, Hoosick Falls***Fusicladium dendriticum* (*Wallr.*) *Fckl.***S. Davis, Boston, Mass.***Clavaria lavendula* *Pk.**Gomphidius maculatus* (*Scop.*) *Fr.**C. pallescens* *Pk.**Inocybe hiulca* *Fr.**Clitocybe brumalis* *Fr.**I. infelix brevipes* *Pk.**C. compressipes* *Pk.**Marasmius varicosus* *Fr.**Clitopilus davisii* *Pk.**Mycena pseudopura* *Cke.**Eccilia watsoni* *Pk.**Naucoria firma* *Pk.**Entoloma griseo-cyaneum* *Fr.**N. sphagnophila* *Pk.**E. sericeum* *Fr.**Nolanea conica* *Pk.**E. variabile* *Pk.**Omphalia pyxidata* (*Bull.*) *Fr.**Galera later. albicolor* *Pk.**Pholiota autumnalis* *Pk.*

J. Dearness, London, Can.

<i>Clasterosporium caricinum Schw.</i>	<i>Marsonia potentillae (Desm.)</i>
<i>Entyloma linariae Schroet.</i>	<i>Microstroma juglandis (Bereng.)</i>
<i>Hypocrella hypoxylon (Pk.)</i>	<i>Peridermium conorum-piceae (Rees)</i>
<i>Isariopsis albo-rosella (Desm.)</i>	<i>Puccinia caricis-asteris Arth.</i>
<i>Leptothyrium punctiforme B. & C.</i>	<i>Sphaerotheca humuli (DC.) Burr.</i>

F. Dobbin, Shushan

<i>Hedeoma hispida Pursh</i>	<i>Stellaria borealis Bigel.</i>
------------------------------	----------------------------------

C. J. Elting, Highland

Centaurea solstitialis L.

C. E. Fairman, Lyndonville

<i>Belonidium glyceriae Pk.</i>	<i>Lophiotrema littorale Speg.</i>
<i>Cantharellus floccosus Schw.</i>	<i>Melanopsamma confertissima (Plow.)</i>
<i>Ciboria luteo-virescens R. & D.</i>	<i>Microsphaera diffusa C. & P.</i>
<i>Diplodia cercidis E. & E.</i>	<i>Ovularia obliqua (Cke.) Oud.</i>
<i>D. hamamelidis Fairm.</i>	<i>Pezizella lanc.-paraphysata Rehm</i>
<i>D. tamariscina Sacc.</i>	<i>Phialea scutula (Pers.) Gill.</i>
<i>Fenestella amorpha E. & E.</i>	<i>Polyporus sulphureus (Bull.) Fr.</i>
<i>Helotium salicellum Fr.</i>	<i>Puccinia epiphylla (L.) Wettst.</i>
<i>Lycogala flavo-fuscum (Ehrh.) Rost.</i>	<i>Stephanoma strigosum Wallr.</i>
<i>Lophiotrema hysterioides (E. & L.)</i>	<i>Trichosporium variabile Pk.</i>

G. C. Fisher, DeFuniak Springs, Fla.

<i>Bovistella floridensis Pk.</i>	<i>Peridermium pyriforme Pk.</i>
-----------------------------------	----------------------------------

W. P. Fraser, Pictou, Can.

<i>Calicium lenticulare (Hoffm.) Ach.</i>	<i>Lycopodium sabinaefolium Willd.</i>
<i>Cenangium populneum (Pers.) Rehm</i>	<i>L. sitchense Rupr.</i>
<i>Dothidella kalmiae (Pk.) Sacc.</i>	<i>Ramularia dubia Riess</i>
<i>Gnomoniella coryli (Batsch) Sacc.</i>	<i>Septogloeum salicinum (Pk.) Sacc.</i>
<i>Venturia pulchella C. & P.</i>	

C. Gaffin, Utica

<i>Volvaria bombycina (Pers.) Fr.</i>

H. Garman, Lexington, Ky.

<i>Pholiota vermiflua Pk.</i>

S. J. Greenfield, Ilion

<i>Panaeolus retirugis Fr.</i>

J. G. Grossenbacher, Geneva

<i>Cryptosporium cerasinum Pk.</i>

M. E. Hard, Kirkwood, Mo.

<i>Laternea columnata Nees</i>

E. T. Harper, Chicago, Ill.

<i>Corticium mutatum</i> <i>Pk.</i>	<i>Myxosporium acerinum</i> <i>Pk.</i>
<i>Diaporthe aucupariae</i> <i>Hazsl.</i>	<i>Phoma lebiseyi</i> <i>Sacc.</i>
<i>Diplodina fusispora</i> <i>Pk.</i>	<i>P. menispermii</i> <i>Pk.</i>
<i>Dothiorella celastri</i> <i>Pk.</i>	<i>P. platysperma</i> <i>Pk.</i>
<i>Fusarium pyrochroum</i> (<i>Desm.</i>) <i>Sacc.</i>	<i>Sphaeropsis simillima</i> <i>Pk.</i>
<i>Helminthosporium macrocarpum</i> <i>Grev.</i>	<i>Stagonospora linearis</i> <i>Pk.</i>
<i>Macrophoma samaricola</i> <i>Sacc.</i>	<i>Stemphylium macrosporoideum</i> (<i>B. & C.</i>)

O. Hill, Boston, Mass.

<i>Agaricus halophilus</i> <i>Pk.</i>	<i>Hypholoma rigidipes</i> <i>Pk.</i>
	<i>Tricholoma subcinereum</i> <i>Pk.</i>

G. T. Howell, Rockville, Ind.

<i>Flammula praecox</i> <i>Pk.</i>	<i>Flammula pulchrifolia</i> <i>Pk.</i>
	<i>Lepiota granosa</i> <i>Morg.</i>

G. Jericho, Albany

<i>Calvatia cyathiformis</i> (<i>Bosc</i>) <i>Morg.</i>

C. E. Jones, Albany

<i>Prunus pumila</i> <i>L.</i>

M. E. Jones, Salt Lake City, Utah

<i>Actinonema rosae</i> (<i>Lib.</i>) <i>Fr.</i>	<i>Linospora brunellae</i> <i>E. & E.</i>
<i>Ascochyta colorata</i> <i>Pk.</i>	<i>Macrosphaera alni ludens</i> <i>Salm.</i>
<i>Cylindrosporium padi cerasinum</i> (<i>Pk.</i>)	<i>M. diffusa</i> <i>C. & P.</i>
<i>C. simile</i> <i>Pk.</i>	<i>Phyllosticta angelicae</i> <i>Sacc.</i>
<i>Dimerosporium collinsii</i> (<i>Schw.</i>)	<i>Physoderma vagans</i> <i>Schroet.</i>
<i>Doassansia alismatis</i> (<i>Nees</i>)	<i>Septoria sacch. occidentalis</i> <i>E. & E.</i>
<i>D. sagittariae</i> (<i>West.</i>)	<i>S. sorbi</i> <i>Lasch</i>
<i>Erysiphe polygoni</i> <i>DC.</i>	<i>S. streptopodis</i> <i>Pk.</i>
	<i>Sphaerotheca humuli</i> (<i>DC.</i>) <i>Burr.</i>

R. Latham, Orient Point

<i>Angelica atropurpurea</i> <i>L.</i>	<i>Iris prismatica</i> <i>Pursh</i>
<i>Arenaria peploides</i> <i>L.</i>	<i>Leontodon nudicaulis</i> (<i>L.</i>) <i>Banks</i>
<i>Aristida gracilis</i> <i>Ell.</i>	<i>Ligusticum scoticum</i> <i>L.</i>
<i>Atriplex pat. littoralis</i> (<i>L.</i>)	<i>Myosotis virginica</i> (<i>L.</i>) <i>B. S. P.</i>
<i>Cerastium viscosum</i> <i>L.</i>	<i>Onopordum acanthium</i> <i>L.</i>
<i>Cirsium spinosissimum</i> (<i>Walt.</i>)	<i>Panicum spretum</i> <i>Schultes</i>
<i>Cyperus nuttallii</i> <i>Eddy</i>	<i>Picris echioides</i> <i>L.</i>
<i>Fimbristylis castanea</i> (<i>Mx.</i>) <i>Vahl</i>	<i>Plantago decipiens</i> <i>Barneoud</i>
<i>Fomes rimosus</i> <i>Berk.</i>	<i>Polygonum littorale</i> <i>Link</i>
<i>Hieracium gronovii</i> <i>Mx.</i>	<i>Rumex pallidus</i> <i>Bigel.</i>
<i>H. scabrum</i> <i>Mx.</i>	<i>Salsola kali</i> <i>L.</i>
<i>Hypericum canadense</i> <i>L.</i>	<i>Silybum marianum</i> (<i>L.</i>) <i>Gaertn.</i>
<i>Hypochaeris radicata</i> <i>L.</i>	<i>Solidago aspera</i> <i>Ait.</i>
<i>Ilex vert. tenuifolia</i> (<i>Torr.</i>)	<i>Strophostyles helvola</i> (<i>L.</i>)
	<i>Tripsacum dactyloides</i> <i>L.</i>

J. Mickleborough, Brooklyn*Myxosporium castaneum* *Pk.***A. J. Miller, Rensselaer**Nuts of *Phytelephas macrocarpa* *R. & P.***G. E. Morris, Waltham, Mass.**

<i>Amanita morrisii</i> <i>Pk.</i>	<i>Eccilia pyrina</i> <i>B. & C.</i>
<i>A. muscaria</i> <i>L.</i>	<i>Entoloma cuspidatum</i> <i>Pk.</i>
<i>A. russuloides</i> <i>Pk.</i>	<i>E. jubatum</i> <i>Fr.</i>
<i>Boletinus grisellus</i> <i>Pk.</i>	<i>E. rhodopolium</i> <i>Fr.</i>
<i>Boletus morrisii</i> <i>Pk.</i>	<i>E. salmoneum</i> <i>Pk.</i>
<i>B. spectabilis</i> <i>Pk.</i>	<i>Geoglossum nigrum</i> <i>Pers.</i>
<i>Calocera cornea</i> <i>Fr.</i>	<i>Hydnum graveolens</i> <i>Delast.</i>
<i>Clitocybe centralis</i> <i>Pk.</i>	<i>H. laevigatum</i> <i>Sw.</i>
<i>C. metachroa</i> <i>Fr.</i>	<i>Lactarius bryophilus</i> <i>Pk.</i>
<i>Coprinus niveus</i> (<i>Pers.</i>) <i>Fr.</i>	<i>Leotia punctipes</i> <i>Pk.</i>
<i>Cortinarius ferrug.-griseus</i> <i>Pk.</i>	<i>Russula serissima</i> <i>Pk.</i>
<i>Eccilia flavida</i> <i>Pk.</i>	<i>Tricholoma piperatum</i> <i>Pk.</i>

W. A. Murrill, New York

<i>Hypholoma boughtoni</i> <i>Pk.</i>	<i>Inocybe infida</i> <i>Pk.</i>
---------------------------------------	----------------------------------

H. S. Paine, Glens Falls*Pholiota duroides* *Pk.***C. R. Pettis, Lake Clear Junction***Peridermium strobili* *Kleb.***H. G. Pierce, Rochester***Salix alba* *L.***E. Riesel, Herkimer***Cuscuta arvensis* *Beyrich***W. H. Ropes, Salem, Mass.**

<i>Lepiota americana</i> <i>Pk.</i>	<i>Lepiota cepaes. lutea</i> (<i>Bolt.</i>)
	<i>Lepiota friesii</i> <i>Lasch.</i>

J. C. Smock, Hudson

<i>Bidens beckii</i> <i>Torr.</i>	<i>Hybanthus concolor</i> (<i>Forst.</i>)
<i>Erythronium albidum</i> <i>Nutt.</i>	<i>Schwalbea americana</i> <i>L.</i>

P. Spaulding, Washington, D. C.*Peridermium strobili* *Kleb.*

E. B. Sterling, Trenton, N. J.*Agaricus eludens* *Pk.**Agaricus magniceps* *Pk.***F. C. Stewart**, Geneva*Erysiphe cichoracearum* *DC.**Hypholoma perplexum* *Pk.**Fomes ribis* (*Schum.*) *Fr.**Microcera coccophila* *Desm.**Phomopsis stewartii* *Pk.***H. L. True**, McConnellsville, O.*Polyporus flavovirens* *B. & R.**Xylaria digitata* (*L.*) *Grev.***B. D. VanBuren & S. H. Burnham**, Albany*Collybia velutipes* (*Curt.*) *Fr.***J. M. VanHook**, Greencastle, Ind.*Hydnum laciniatum* *Leers***H. Wardell**, Middleburg*Lappula virginiana* (*L.*) *Greene***H. L. Wells**, New Haven, Conn.*Agaricus rodmani* *Pk.***F. B. Wheeler**, Syracuse*Morchella crispa* *Karst.**Morchella rimosipes* *DC.**Pholiota aurivella* *Batsch***H. H. Whetzel**, Ithaca*Ascochyta solani-nigri* *Diedicke***T. E. Wilcox**, Washington, D. C.*Amanita flavorubescens* *Atk.***D. B. Young**, Albany*Amanitopsis vaginata* (*Bull.*) *Roze**Entoloma salmoneum* *Pk.**Boletus albus* *Pk.**Lactarius deliciosus* *Fr.**B. piperatus* *Bull.**L. oculatus* (*Pk.*) *Burl.**Eccilia atrides* *Lasch.**Russula fragilis* (*Pers.*) *Fr.**Sporotrichum larvatum* *Pk.*

SPECIES NOT BEFORE REPORTED

Ascochyta solani-nigri Diedicke

Living leaves of egg plant, *Solanum melongena* L.
Ithaca. October. H. H. Whetzel.

Agropyrum tenerum Vasey

Brownsville, Jefferson co. and Adirondack mountains. June and July. Formerly confused with *Agropyrum violaceum* Lange.

Belonidium glyceriae n. sp.

Receptacle 1-1.5 mm broad, gregarious, sessile, plane or convex, glabrous or merely papillate on the under side, pale yellow; asci subclavate or subfusiform, obtuse, 120-130 x 14-18 μ ; spores oblong or subcylindric, straight or slightly curved, 3-septate, often 4-nucleate, crowded or biseriate, 35-40 x 4-5 μ , paraphyses filiform.

Dead culms of *Glyceria nervata* (Willd.) Trin. Lyndonville, Orleans co. June. C. E. Fairman.

Receptaculum 1-1.5 mm latum, gregarium, sessile, planum convexumve, extus glabrum seu papillatum, flavidum; asci subclavati vel subfusiformes, obtusi, 120-130 x 14-18 μ ; sporae oblongae vel subcylindratae, rectae vel leviter curvae, 3-septatae, saepe 4-nucleatae, confertae vel distichae, 35-40 x 4-5 μ , paraphyses filiformes.

Biatora cupreo-rosella (Nyl.) Tuckerm.

Limestone rocks. Pine Island, Orange co. November. C. F. Austin.

Bidens tenuisecta Gray

Field near Rochester. August. Miss F. Beckwith. Probably a recent introduction from the West.

Boletus viridarius Frost

Grassy places near pine trees. Poughkeepsie. September and October. Miss H. L. Palliser. For description of this species see article on "Edible fungi" in another part of this report.

Bromus altissimus Pursh

Rathbone, Steuben co. and North Greenbush, Rensselaer co. August to October. Formerly confused with *Bromus ciliatus* L.

***Cardamine douglasii* (Torr.) Britton**

Niagara Falls and Syracuse. May. Formerly referred to *Cardamine rhomboidea purpurea* Torr., but now recognized as a distinct species.

***Carduus crispus* L.**

Fields. Helderberg mountains. October. S. H. Burnham. A recently introduced plant, very spiny but beautiful.

***Carex bebbii* Olney**

Common. Formerly considered a variety of *Carex tribuloides* Wahl., but now recognized as a distinct species.

***Carex crawfordii* Fern.**

Common. Previously known as *Carex scoparia minor* Boott, but raised to specific rank in the New Manual.

***Chaenactis stevioides* H. & A.**

Newly seeded lawn. Rochester. Miss F. Beckwith. Probably a recent introduction from the West.

***Ciboria luteo-virescens* R. & D.**

On petioles of fallen maple leaves. Lyndonville. C. E. Fairman.

***Clitocybe candida* Bres.**

Woods. West Fort Ann, Washington co. October. S. H. Burnham. The pileus in these specimens is not a pure white as might be inferred from the specific name, but is tinged in the center with yellowish or grayish brown hues. It is also sometimes eccentric.

***Cortinarius subsalmoneus* Kauffm. Ms.**

Woods. Hague, Warren co. September. S. H. Burnham. The full description of this species has not yet been published, but the specimens agree with those characters published in the *Key to the Species of Cortinarius* and in the author's manuscript description.

***Crataegus brevipes* n. sp.**

Leaves ovate or broadly ovate, acute, rounded or broadly cuneate at the base, with 2-3 very slight broad lobes each side or scarcely lobed, with marginal teeth short, broad and blunt, glabrous except

a few hairs on the upper surface of the midrib near the base, slightly bronze tinged when unfolding, soon green or yellowish green, paler beneath, becoming darker green and firm, those on vigorous shoots larger, subcoriaceous, more distinctly lobed, and broadly rounded or subtruncate at the base, petioles short, 6-12 mm long, slightly margined at the top, nearly or quite glandless.

Flowers 5-10 in a cluster, 1.6-2.4 mm broad, commonly on simple glabrous pedicels 6-12 mm long, calyx lobes irregular, often abruptly narrowed toward the reddish apex, entire or with few marginal glands, slightly hairy inside; stamens 8-10, anthers pink; styles 3-4.

Fruit erect, globose or depressed globose, 1-1.4 cm long, 1.2-1.4 cm broad, angular, scarcely or not at all pruinose, 3-8 in a cluster, supported on short glabrous pedicels, dull red or blotched with green, nutlets 3-4, 7-8 mm long.

A shrub 2-3 m tall, with wide spreading branches armed with stout, nearly straight spines 2.5-4 cm long. Flowers the last week in May. Fruit ripe the latter part of September.

Rocky hilly places. Corning, Steuben co. The species evidently belongs to the Pruinosa group, though the fruit is not distinctly pruinose. The specific name has reference to the short pedicels, by which character the species is distinguished from all others of this group known to me.

Folia ovata vel late ovata, acuta, basi rotundata vel late cuneata, utriusque 2-3 lobata, seu vix lobata, dentibus brevibus, latis, obtusis, margine serrata, glabra, nisi pilis paucis in venis ad basem, juventute leviter rufobrunnea, mox viridia, infra pallidiora, in maturitate virescentiora et subcoriacea, petiolae 6-12 mm longae, ad apicem leviter marginatae fere eglandulosae.

Flores 5-10 in corymbo, 1.6-2.4 cm latae, in pedicellis glabris, vulgo simplicibus, 6-12 mm longis, calicis lobi irregulares, saepe ad apicem abrupte angustati et rubri, integri vel glandis paucis, intra leviter hirti, stamines 8-10, antherae rosaceae; styles 3-4.

Poma 1-1.4 cm longa, 1.2-1.4 lata, angularia, epruinosa, brevibus glabris erectis pedicellis suffulta, sanguinea, nuculae 3-4, 7-8 mm longae.

***Diplocladium penicilloides* Sacc.**

Decaying specimens of *Polyporus resinosus* (Schrad.) Fr. Helderberg mountains. May. S. H. Burnham.

Diplodia cercidis E. & E.

Dead branches of cultivated *Cercis japonica* Sieb. Lyndonville. July. C. E. Fairman.

Diplodia hamamelidis n. sp. Fairm. in litt.

Perithecia gregarious, minute, depressed globose, black, at first covered by the epidermis, then erumpent; spores at first colorless, then colored, for a long time continuous, finally uniseptate, $20-28 \times 10-12 \mu$.

Dead branches of witch hazel, *Hamamelis virginiana* L. Lyndonville. September. C. E. Fairman.

Perithecia gregaria, minuta, depresso-globosa, nigra, primus epidermide tecta, deinde erumpentia; sporae primus hyalinae, deinde coloratae, diu continuae, denique uniseptatae, $20-28 \times 10-12 \mu$.

Diplodia tamariscina Sacc.

Dead branches of cultivated *Tamarix parviflora* DC. Lyndonville. July. C. E. Fairman.

Discina leucoxantha Bres.

Ground, under beech trees. Altamont, Albany co. May.

Dothiorella divergens n. sp.

Clusters of perithecia 1-3 mm broad, seated on or immersed in a black stroma, suborbicular or elliptic, erumpent, surrounded by the ruptured epidermis; perithecia irregular, unequal, submembranous, black, pallid within; spores oblong, obtuse, straight or slightly curved, sometimes uninucleate, hyaline rarely becoming greenish or yellowish, $20-30 \times 9-10 \mu$.

Dead branches of apple tree, *Pyrus malus* L. Menands, Albany co. May.

This species differs from *Dothiorella mali* E. & E. and *D. pyrenophora* Karst. & Sacc. in its much larger spores. It diverges from the generic character in sometimes having spores slightly tinged with green or yellow.

Caespites perithecorum 1-3 mm lati, insidentes stromate nigro, suborbiculares seu ellipsoidei, erumpentes, epidermide rupta circumdati; perithecia irregularia, inaequalia, submembranacea, atra, intus pallida; sporae oblongae, obtusae, rectae, vel leviter curvae, aliquando uninucleatae, hyalinae, rare viridescentes seu flavescentes, $20-30 \times 8-10 \mu$.

***Epipactis tessellata* (Lodd.) Eaton**

Woods. Gansevoort, Saratoga co. August.

***Fenestella amorphia* E. & E.**

Dead branches of hickory. Lyndonville. July. C. E. Fairman.

***Geum flavum* (Port.) Bickn.**

Greenburg, Westchester co. E. C. Howe. Shushan, Washington co. September. S. H. Burnham.

✓ *Hypholoma boughtoni* n. sp.

PLATE II, FIG. I-7

Pileus fleshy, thin except in the center, broadly convex or sub-hemispheric, rarely with a slight umbo, glabrous or slightly fibrillose, often concentrically and areolately cracking, pale reddish brown or grayish brown, flesh whitish, taste disagreeable; lamellae unequal, moderately close, adnate, purplish brown, seal brown or blackish, obscurely spotted, whitish on the edge; stem equal, floccosely fibrillose, striate at the top, hollow, white or whitish; spores black on white paper, broadly elliptic, apiculate, $10-12 \times 7-8 \mu$.

Pileus 2.5-7 cm broad; stem 2.5-6 cm long, 4-10 mm thick.

Ground in woods and in open places. Near Pittsford, Monroe co. and at Menands. August. F. S. Boughton and C. H. Peck.

This species is closely allied to *Hypholoma velutinum* (Pers.) Fr. from which it may be separated by its dry, not hygrophanous, pileus, its whitish flesh and stem, the absence of cystidia and the larger spores. The spore print of both this and *Hypholoma rigidipes* Pk. is black on white paper. This would indicate a close relationship to the *Melanosporae*, not only of these two species, but probably also of the closely related species *H. velutinum* (Pers.) Fr. and *H. lacrymabundum* Fr.

Pileus carnosulus, centro excepto, late convexus vel subhemisphaericus, rare subumbonatus, glaber vel subglaber, saepe rimosus, rufo-brunneus vel griseo-brunneus, carne albida, sapore ingrato; lamellae inaequales, subconfertae, adnatae, obscure maculatae, purpureo-brunneae, atro-brunneae vel nigrescentes, acie albida; stipes aequalis, floccoso-fibrillosus, ad apicem striatus, cavus, albidus; sporae in fundamento candido atrae, late ellipsoideae, apiculatae, $10-12 \times 7-8 \mu$.

Hypholoma rigidipes n. sp.

PLATE III, FIG. I-6

Pileus fleshy, thin, convex or broadly convex, dry, fibrillose-squamulose, tawny brown (raw umber), often reddish in the center, flesh whitish, odor slight or none, taste mild; lamellae narrow, close, slightly sinuate, adnexed, brownish red becoming black or purplish black with age; stem slender, rigid, equal, hollow, fibrillose squamulose, colored like the pileus or a little paler; spores subellipsoid, apiculate, $10-12 \times 6-8 \mu$.

Pileus 2.5-5 cm broad; stem 5-10 cm long, 4-6 mm thick.

Gregarious. Damp places under tall herbs. North River, Warren co. September.

This is closely related to *Hypholoma lacrymabundum* Fr. from which it may be separated by its smaller size, gregarious mode of growth, slender, rigid, equal, darker colored stem, larger spores and slight evanescent veil.

Pileus carnosulus, convexus vel late convexus, siccus, fibrilloso-squamulosus, umbrinus, saepe in centro rubescens, carne albida, sapore mite; lamellae angustae, confertae, adnexae leviter sinuatae, rufo-brunneae, deinde purpureo-atrae vel nigrescentes; stipes gracilis, rigidus, aequalis, cavus, fibrilloso-squamulosus, pileo in colore similis; sporae subellipsoideae, apiculatae, $10-12 \times 6-8 \mu$.

Juncus brachycephalus (Engelm.) Buchen.

Jamesville, Onondaga co. and Sevey, St Lawrence co. July and August. Formerly reported as a variety of *Juncus canadensis* Gay.

Juncus brevicaudatus (Engelm.) Fern.

West Albany, Sand Lake and Adirondack mountains. August and September. Formerly reported as a variety of *Juncus canadensis* Gay.

Juncus secundus Beauv.

Blue Mountain Lake, Hamilton co. August. Reported as a variety of *Juncus tenuis* Willd.

Juniperus horizontalis Moench

Bergen swamp, Genesee co. Formerly reported as a variety of *Juniperus sabina* L.

Leontodon nudicaulis (L.) Banks

Orient Point, Suffolk co. September. R. Latham.

Leskea gracilescens Hedw.

Trunks of trees. Kingston. July. Mrs M. E. Williams.

Ligusticum scoticum L.

Orient Point. September. R. Latham.

Lophiotrema hysterioides E. & E.

Decorticated twigs and branches lying on the ground in woods. Lyndonville. September. C. E. Fairman.

Lophiotrema littorale Speg.

Dead branches of willow. Lyndonville. May. C. E. Fairman.

✓ **Marasmius alienus** n. sp.

Pileus thin, tough, convex, subpruinose, dry, pallid or pale buff, with a thin straight margin; lamellae subarcuate, distant, slightly decurrent, creamy yellow, becoming brownish in drying; stem slender, firm, hollow, subpruinose, pallid; spores oblong or narrowly elliptic, $8-10 \times 4-5 \mu$.

Pileus 6-12 mm broad; stem 2.5-5 cm long, .5-1 mm thick.

Mossy prostrate trunks of trees in woods. Fine, St Lawrence co. August.

This species belongs to section 3, subsection 2, of Professor Morgan's *Synopsis of North American Species of Marasmius*.

Pileus tenuis, lentus, convexus, subpruinosis, siccus, pallidus vel subluteolus, margine tenue, recto; lamellae subarcuatae, distantes, leviter decurrentes, cremeae, in siccitate brunnescentes; stipes gracilis, firmus, fistulosus, subpruinosis, pallidus; sporae oblongae vel anguste ellipsoideae, $8-10 \times 4-5 \mu$.

Melanopsamma confertissima (Plowr.) Sacc.

Dead branches of spice bush, *Benzoin aestivale* (L.) Nees. Lyndonville. C. E. Fairman.

Microcera coccophila Desm.

Parasitic on San José scale infesting living branches of apple trees. Hicksville, Nassau co. October. F. C. Stewart. A welcome enemy to a very unwelcome foe to fruit trees and shrubs.

Midotis irregularis (Schw.) Cke.

On decaying wood. Indian Lake, Hamilton co. October.

Monolepis nuttalliana (R. & S.) Wats.

Rochester. Miss F. Beckwith. September. An interesting member of the Goosefoot family. It has probably been recently introduced from the western part of the country.

Morchella crispa Karst.

Near Syracuse. May. F. B. Wheeler. It resembles *Morchella conica* Pers. but may be distinguished from it by the more irregular tortuous ribs of the cap, the chinks at the base of the stem and the longer spores.

Morchella rimosipes DC.

Near Syracuse. May. F. B. Wheeler. Probably both this and the preceding species of morel are edible, but as I have had no opportunity of making a personal test of their edible quality they are not here recorded as such.

Naias gracillima (A. Br.) Magn.

Water holes near West Albany. September. Formerly reported as a variety of *Naias indica* Willd. but now considered a distinct species.

Nardia crenulata (Sm.) Lindb.

Near Calamity pond, Essex co. August. Miss A. Lorenz.

Nardia hyalina (Lyell) Carr.

Banks of Marcy brook, Essex co. July. Miss A. Lorenz.

Panicum implicatum Scribn.

Albany; Machias, Cattaraugus co. and Adirondack mountains. July. Formerly confused with *Panicum pubescens* Lam. and *P. lanuginosum* Ell.

Panicum oricola H. & C.

Manor and Riverhead, Suffolk co. and Fulton Chain, Herkimer co. July and August. Formerly confused with *Panicum dichotomum* L. and *P. atlanticum* Nash.

***Panicum spretum* Schultes**

Near Albany; Riverhead and Orient Point, Suffolk co. and Whitehall, Washington co. July. Formerly confused with *Panicum dichotomum* L.

***Peridermium strobil* Kleb.**

Seedling white pines, *Pinus strobus* L. Lake Clear Junction, Franklin co. October. Perley Spaulding and C. R. Pettis. Our specimens are immature.

This parasitic fungus is destructive to white pine trees. It is dimorphic. *Cronartium ribicola* Dietr. is a form which develops on leaves of currant bushes. Its spores are capable of infecting white pine trees and reproducing the pine rust, *Peridermium strobil*, in them. To prevent this it is important that currant and gooseberry bushes whose leaves are attacked by the *Cronartium* should be destroyed at once.

***Pezizella lanceolato-paraphysata* Rehm**

Dead stems of cultivated *Spiraea filipendula* L. Lyndonville. June. C. E. Fairman.

***Phaeopezia fuscocarpa* (E. & H.) Sacc.**

Decaying wood. Kasoag, Oswego co. July.

***Pholiota aurivella* Batsch**

Decaying wood of maple. Near Syracuse. October. F. B. Wheeler.

***Phomopsis stewartii* n. sp.**

Perithecia gregarious, commonly occupying grayish or brown spots, thin, subcutaneous, at length erumpent, depressed, minute, $\frac{1}{3}$ – $\frac{1}{2}$ mm broad, black; spores of two kinds, first, filiform, curved, flexuous or uncinat, hyaline, 16 – 25×1 – 1.5μ , second, oblong or subfusiform, hyaline, commonly binucleate, 8 – 12×2 – 3μ ; sporophores slender, equal to or shorter than the spores.

On stems of *Cosmos bipinnatus* Cav. Garden of Agricultural Experiment Station, Geneva, Ontario co. October. F. C. Stewart.

Perithecia gregaria, maculas griseas seu brunneas vulgo occupantia, tenua, subcutanea, deinde erumpentia, depressa, minuta, $\frac{1}{3}$ – $\frac{1}{2}$ mm lata, nigra; sporae dimorphae, primum, filiformes, cur-

vatae, flexuosae hamataeve, hyalinae, 16-25 x 1-1.5 μ , secundum, oblongae vel subfusiformes, hyalinae, vulgo binucleatae 8-12 x 2-3 μ , sporophores graciles, sporis aequales vel breviores.

Picris echinoides L.

Orient Point. September. R. Latham.

Potamogeton richardsoni (Benn.) Rydb.

Lake Champlain and Oneida lake. August. Formerly reported as a variety of *Potamogeton perfoliatus* L. but now classed as a distinct species.

✓ **Psilocybe nigrella n. sp.**

PLATE III, FIG. 7-II

Pileus thin, broadly convex or nearly plane, slightly umbonate, hygrophanous, seal brown and shining when moist, even and obscurely striate on the margin, raw umber or mummy brown when the moisture has escaped; lamellae thin, rather close, rounded behind, adnexed, purple brown or seal brown, whitish on the edge; stem firm, rigid, equal, stuffed with a slender white pith, silky fibrillose, whitish; spores dark purplish brown, almost black, ellipsoid, 10-12 x 6-8 μ .

Pileus 2.5-4 cm broad; stem 3.5-7 cm long, 2-4 mm thick.

Damp mossy ground in swamps. Karner, Albany co. October.

Pileus tenuis, late convexus subplanusve, leviter umbonatus, hygrophanous, atrobrunneus, nitidus, levis, margineque obscure striatus in conditione uda, umbrinus in siccitate; lamellae tenues, subconfertae, adnexae, atrobrunneae, acie albidae; stipes firmus, rigidus, aequalis, medulla alba farctus, sericeo-fibrillosus, albidus; sporae purpureo-brunneae vel subatrae, ellipsoideae, 10-12 x 6-8 μ .

Puccinia epiphylla (L.) Wettst.

Living leaves of low spear grass, *Poa annua* L. Lyndonville. September. C. E. Fairman.

Ribes triste albinervium (Mx.) Fern.

Colton hill swamp. Fine. August.

This is closely related to *Ribes vulgare* Lam., the common garden currant, from which it may be separated by its more straggling, partly decumbent habit and by the glands on the pedicels.

The variety differs from the typical form in having the leaves glabrous on the lower surface.

Rubia tinctorum L.

Near Smiths Basin, Washington co. October. S. H. Burnham.

Introduced and formerly cultivated for its roots, which yield a coloring matter suitable for dyeing. It has persisted several years in the locality cited.

Rumex pallidus Bigel.

Orient Point. July. R. Latham.

Rubus andrewsianus Blanch.

Sandy soil. Islip, Suffolk co. Formerly considered a small form of *Rubus villosus frondosus* Bigel.

Rubus permixtus Blanch.

Light soil in pastures. North Elba, Essex co. July. Formerly referred to *Rubus procumbens* Muhl.

Rubus recurvans Blanch.

Pine Plains, Dutchess co. and Snyders Corners, Rensselaer co. July and August. Formerly considered a variety of *Rubus villosus* Ait. of the older botanies.

✓ **Septoria sedicola n. sp.**

Spots orbicular, 4-8 mm broad, usually one or two on a leaf, at first definite, depressed and without discoloration of the surrounding leaf tissue, at length convex above, concave beneath, the surrounding part of the leaf becoming yellowish, thin and flaccid, finally the whole leaf dying; perithecia numerous, minute, amphigenous, black; spores filiform, straight, curved or flexuous, enucleate, 20-40 μ long, 1-1.5 μ thick.

Living leaves of live-for-ever, *Sedum purpureum* Tausch. (*Sedum telephium* L. Man. ed. 6.) Fine. August.

This fungus is closely allied to *Septoria sedi* West. from which I have separated it because of its amphigenous perithecia and its enucleate spores. The host plant is very tenacious of life and on that account a very undesirable weed though it spreads slowly. This parasite is injurious to it and tends to keep it in check and may therefore be considered a beneficial fungus.

Maculae orbiculares, 4-8 mm latae, vulgo in ullo folio una duove, primus definitae, depressae, in partibus folii circumdantibus, absque decoloratione, deinde convexae supra, concavae infra, foliis flavescenscentibus, tenuibus, flaccidis, moribundis; perithecia numerosa, minuta, amphigena, nigra; sporae filiformes, rectae, curvatae flexuosaeve, enucleatae, 20-40 x 1-1.5 μ .

***Solidago aspera* Ait.**

Orient Point. September. R. Latham.

***Sparganium americanum* Nutt.**

Lakes and ponds. Sand Lake, Rensselaer co. July. Formerly regarded as *Sparganium simplex nuttallii* Engelm.

***Sparganium angustifolium* Mx.**

Lake Placid, Essex co. Formerly recorded as *Sparganium simplex angustifolium* (Mx.) Engelm.

***Sparganium diversifolium* Graebn.**

Shushan. September. S. H. Burnham.

***Stachys sieboldii* Miq.**

Along the railroad near Whitehall. September. S. H. Burnham.

This is sometimes designated as *Stachys tuberifera* Naud., a name suggested by its tuberous edible roots. It bears the common names knot root, Chinese artichoke and Japan artichoke.

***Stephanoma strigosum* Wallr.**

In woods. Lyndonville. August. C. E. Fairman.

This fungus is parasitic on *Lachnea hemispherica* Wigg. In the generic and specific descriptions some of the characters of the host plant are confusingly incorporated as if they belonged to the parasite.

***Sterigmatocystis ochracea* (Wilh.) VanTigh.**

On diseased gladiolus bulbs and other vegetable matter kept under a bell glass in the office of the State Entomologist. Albany. April and May. J. B. Bartlett.

✓ *Trametes merisma* n. sp.

Pileus coriaceous, fibrous, tough, commonly deeply divided into several pileoli, uneven, tuberculose, colliculose or diminutively proliferous, subpubescent, white or whitish, flesh pure white, the margin obtuse, sterile beneath; pores minute, 2-3 in a millimeter, developing from the center toward the margin, white, the edge of the dissepiments at first obtuse; stem like base short or none; spores not seen.

Pileus 2.5-7 cm broad.

Decaying prostrate trunks of beech trees, *Fagus grandifolia* Ehrh. Fine. August.

This singular fungus sometimes develops from the lower surface of the trunk, in which case a tubercle first appears and the pendent pileus develops from it and is centrally attached to it by the apex. The context of the pileus is similar to that of species of *Polystictus*, but the character of the pores indicates a closer connection with *Trametes*.

Pileus coriaceous, fibrosus, lentus, vulgo in pileolos paucos profunde divisus, asper tuberculosus colliculosus vel leviter proliferus, subpubescens, albus albidusve, carne candida, margine obtuso, infra sterile; pori minuti, .3-.5 mm lati, a centro ad marginem patescentes, albi, dissepimentis obtusis, stipes brevis vel nullus.

✓ *Trichosporium variabile* n. sp.

Widely effused, forming thin indefinite blackish patches; hyphae prostrate or suberect, simple or branched, continuous or rarely septate, 4-5 μ in diameter, brown by transmitted light or partly hyaline; spores varying from globose to oblong, colored, 6-10 μ in diameter or 8-12 x 6-8 μ .

On building paper kept in rolls under shelter. Lyndonville. September and November. C. E. Fairman.

The species is remarkable for the variability in the size and shape of the spores. They are intermingled, but the oblong spores are more numerous than the globose. It differs from *Trichosporium chartaceum* (Pers.) Sacc. in its much larger spores.

Late effusum, stratum tenue indeterminatum nigrescens formans; hyphae repentes suberectaeve, simplices ramosaeve, continuae vel leviter septatae, 4-5 μ crassae, fuscae seu partim hyalinae; sporae fuscae, globosae, 6-10 μ latae, vel oblongae, 8-12 x 6-8 μ .

Verticillium rexianum Sacc.

Parasitic on *Arcyria cinerea* (Bull.) Pers. Fine. August.

Volvaria volvacea (Bull.) Fr.

Pittsford. August. F. S. Boughton. This is a white form with the silky fibrils of the pileus paler than in the typical form.

Zizania palustris L.

Shores of Lake Champlain near Whitehall and Dresden. August and September. This grass was formerly confused with *Zizania aquatica* L. but is now separated as a distinct species, distinguished by its broader leaves.

REMARKS AND OBSERVATIONS

Cardamine bulbosa (Schreb.) B. S. P.

This name takes the place in the New Manual of *Cardamine rhomboidea* DC. in Gray's Manual, ed. 6. Fine flowering specimens of it were collected in May near Little's pond, Albany co.

Centaurea solstitialis L.

This recently introduced plant is apparently spreading, specimens having been received the past season from Highland, Ulster co., where it was collected by C. J. Elting, and from the southern part of Cayuga co., collected by A. D. Baker.

Cerastium viscosum L.

Orient Point. May. R. Latham. This is a rare species in our State.

Crataegus verecunda gonocarpa n. var.

Leaves thin, elliptic, oval or suborbicular, obtuse or acutish, rounded at the base, yellowish green, green with age, becoming glabrous except a few scattered hairs on the upper surface, not at all or only slightly broadly lobed above the middle, the margins often curved upward, petioles 4-10 mm long, slightly margined above, glabrous or with few hairs in the furrow, with few or no glands.

Flowers 5-10 in a cluster, 12-14 mm broad, on short mostly simple pedicels less than an inch long, calyx lobes linear, glabrous, subentire; stamens 1-7, anthers whitish; styles 2-3.

Fruit erect or nearly so, compressed or obtusely 3-angled, dark red when ripe, with numerous minute yellowish lenticels, 10-12 mm long, 7-12 mm broad, compressed fruits about 7 mm in the narrow diameter, 12 mm in the broad diameter, flesh greenish yellow, hard and dry, nutlets 2-3, 7-9 mm long.

Rocky hillside. Corning. May 29, September 17 and 21.

A shrub 2-3 m tall with numerous spreading branches armed with curved spines 2.5-4 cm long, commonly pointing toward the base. The leaves on vigorous shoots are larger than the others, nearly orbicular and more distinctly lobed. The characters which specially distinguish this from the typical plant are its more entire elliptic or suborbicular leaves, its shorter pedicels and its compressed or bluntly angular fruit.

A planta typica differt in foliis magis integris, ellipticis vel sub-orbicularibus, pedicellis brevioribus et fructibus compressis vel obtuse triquetris.

Epilobium densum Raf.

Waste places. Fine. August. The name here used takes the place of *Epilobium lineare* Muhl. used in *Gray's Manual*, ed. 6.

Erythronium albidum Nutt.

This plant formerly grew in the vicinity of Albany but it long ago disappeared from this region. A specimen of it has been contributed by Prof. J. C. Smock, which was collected many years ago and has "Albany" on the label. A specimen in Beck herbarium is labeled "Wet meadows, Albany."

Exoascus pruni Fckl.

This parasitic fungus, which causes the enlargement of the fruit known as "bladder plums," was very prevalent about Rossie, St Lawrence co., in June. Many trees of both the wild black plum, *Prunus nigra* Ait., and the wild red plum, *Prunus americana* Marsh., had scarcely a sound plum on them.

Galium erectum Huds.

This introduced species of bedstraw is abundant in pastures and along roadsides near Hudson. It spreads by subterranean root-stocks and threatens to be a pernicious weed.

Hedeoma hispida Pursh

Two stations are now known in the State for this rare plant, Little Falls and Shushan.

Ilex verticillata tenuifolia (Fern.) Wats.

Orient Point. July. R. Latham.

Lactuca scariola integrata G. & G.

In the *New Manual* this name designates the wild lettuce previously referred to *Lactuca virosa*. This lettuce has now become very common in and around many cities and villages in the State.

Laportea canadensis L.

There are two forms of this nettle. One is common in moist or wet places. It has a slender stem, thin leaves on long slender petioles and usually bears a terminal cluster of pistillate flowers only.

The other is rare, grows in dry soil or upland either in woods or open places, has a stouter stem, thicker leaves on shorter petioles and frequently bears staminate flowers in the axils of most of the leaves, either with or without a terminal cluster of pistillate flowers. This form was found by the roadside at Fine and in woods near Castorland, Lewis co.

Listera australis Lindl.

A single plant was found in a large swamp near Fine. August.

Marasmius oreades Fr.

A variety with the pileus white or whitish occurs in grassy ground at Rossie. September.

✓ *Omphalia rugosodisca levidisca* n. var.

Decaying wood. Fine. August. This differs from the typical form only in having the center of the pileus even.

Pileus in centro levis.

Peridermium consimile A. & K.

Leaves of black spruce, *Picea mariana* (Mill.) B. S. P. Fine. August.

Polyporus giganteus (Pers.) Fr.

About old stumps in woods. Fine. August. This species forms large clusters of pilei which are at first whitish or pale grayish brown, but they become brown or blackish brown in age or in drying. The minute white pores when fresh assume a blackish color where bruised and sometimes become black in drying.

Prunus pumila L.

Pulaski, Oswego co. August. C. E. Jones. The plants growing in sandy soil northwest of Albany and formerly referred to *Prunus pumila* are now referred to *Prunus cuneata*

Raf., which species is characterized by its more erect mode of growth.

***Pyrus coronaria* L.**

The leaves of the American crabapple are quite variable. In one form they are gradually narrowed toward the acute apex but broad at or near the base and often somewhat lobed; in the other they are more or less oblong or elliptic and barely acute at the apex.

***Pyrus melanocarpa* (Mx.) Willd.**

Fruiting specimens of this species were collected at the same time from shrubs on opposite sides of a path in a swamp near Fine. The shrub on one side of the path had black fruit, on the other, dark red.

***Solanum nigrum* L.**

Although the fruit of this plant is reputed poisonous, nevertheless in some places it is used in making pies. The plant is even cultivated for its fruit. A form bearing very large fine fruit was observed in a garden at Rossie and the proprietor assured me that he used the fruit for food. Cooking appears to destroy its deleterious qualities. The cultivated form is locally known as "garden huckleberry."

***Solidago squarrosa ramosa* n. var.**

Corning. September. This differs from the ordinary form in developing a pyramidal panicle of flowers at the top of the stem. The branches are 2.5-10 cm long, gradually diminishing in length from the base to the top of the panicle. The ray flowers are 8-11, disk flowers 9-14. Leaves more narrow than in the common form.

Panícula pyramidata, ramis 2.5-10 cm longis, flores marginis 8-11, flores disci 9-14, folia angustiora.

***Schwalbea americana* L.**

In the *New Manual* this plant is said to grow in wet sandy soil near the coast. In Beck's *Botany* it is credited to sandy plains near Albany. In Paine's *Catalogue of Oncida County Plants* it is reported as occurring near Center (Karner) station between Albany and Schenectady. A specimen has been contributed to the herbarium by Professor Smock that was credited to Albany and probably collected in or near the locality observed by the author

of Paine's *Catalogue*. In the Beck herbarium there are specimens credited to Albany.

***Thalictrum confine* Fern.**

Rossie. June. Specimens collected near Port Henry and referred to *Thalictrum purpurascens* L. belong here.

***Thalictrum revolutum* DC.**

This name is used in the *New Manual* to designate the plant formerly referred to *Thalictrum purpurascens ceriferum* Aust. and the glandular leaved form of *Thalictrum purpurascens* L.

***Viola sororia* Willd.**

A white or whitish flowered form of this species was found near Rochester in May by Miss F. Beckwith and specimens were contributed by her to the herbarium.

EDIBLE FUNGI

***Clitocybe multiceps* Pk.**

MANY CAP CLITOCYBE

PLATE II7, FIG. 7-9

Pileus fleshy, firm, convex, slightly moist in wet weather, whitish, grayish or yellowish gray, flesh white, taste mild; lamellae close, adnate or slightly decurrent, whitish; stems densely cespitose, equal or slightly thickened at the base, solid or stuffed, firm, slightly pruinose at the top, whitish; spores globose, .0002-.0003 of an inch in diameter (5-8 μ).

The many cap clitocybe is quite constantly tufted in its mode of growth. The tufts may be composed of two or three or many individuals. When there are many individuals in a tuft the caps are generally irregular because closely crowded against each other in their growth. The surface is smooth but sometimes slightly silky and brownish in the center. The color is whitish, grayish or yellowish gray, but the flesh is pure white. The gills are white, closely placed, with intervening short ones, the longest ones reaching the stem and broadly connecting with it or slightly decurrent on it. The stems are stout, nearly equal in diameter in every part.

smooth, solid, white or whitish. They are crowded or even attached to each other at the base.

They may appear at any time from June to October if the weather is sufficiently rainy. The taste, though not acrid, is sometimes slightly disagreeable in the raw state, and unless thoroughly cooked the disagreeable flavor may not be wholly dispelled in preparing the caps for the table. This has given rise to different opinions concerning its edibility. One correspondent declares that he considers it one of the best mushrooms. Another thinks it unfit to eat. My first trials of it were not satisfactory. More recent ones lead me to place it among our edible species though it is scarcely to be considered first-class.

***Lactarius aquifluus* Pk.**

WATERY MILK LACTARIUS

PLATE II8, FIG. 1-6

Pileus fleshy, fragile, convex or nearly plane, at length centrally depressed, sometimes with a small umbo, glabrous or slightly and minutely tomentose, burnt sienna red when young and moist, paler grayish buff or subochraceous when dry, flesh colored nearly like the pileus, milk watery, taste mild or tardily acrid; lamellae thin, close, adnate or slightly decurrent, yellowish; stem equal or slightly tapering upward, glabrous or subpruinose, hollow, paler than the pileus; spores subglobose, .0003-.00035 of an inch in diameter (8-9 μ).

The watery milk lactarius grows in mossy swamps or wet places, rarely as a short stem variety, *Lactarius aquifluus brevissimus* Pk., in black muck soil in old roads in woods. The plants are generally gregarious but sometimes tufted. The cap is 2-4 inches broad, the stem 1-4 inches long and 4-8 lines thick. It is moist or subhygrophanous in wet weather and even in dry weather when growing in wet places.

The color of the cap is at first yellowish red, but this soon changes to a grayish or pale ochraceous color as the moisture escapes. The flesh is colored similar to the pileus. The milk is scant and watery in appearance. The taste is mild or slowly and slightly acrid. The odor in the fresh plant is weak but agreeable. It becomes stronger in the dried plant and persists a long time. It is not always entirely destroyed even in cooking. It resembles the odor of melilot and is similar to that of *Lactarius glyciosmus* Fr. and *Lactarius camphoratus* (Bull.) Fr. The

gills have a pale creamy yellow color and become pruinose dusted by the spores in the dried plant. They are at first broadly attached to the stem but in specimens having the pileus centrally depressed they become slightly decurrent. The stem is nearly or quite smooth, hollow and colored like, but a little paler than the cap. It is generally about equal to the diameter of the cap in length. In the upland form it is shorter.

The species is closely related to *Lactarius helvus* Fr. of Europe, which is said by Fries to occur in a degenerate form in swamps and to have a rimose cap and watery milk. If we admit that Fries was correct in considering his watery milk *lactarius* a degenerate form of his typical *Lactarius helvus* with white milk, it still remains doubtful if our plant is the same as his, as some have claimed. The reasons for considering it a distinct species are two. First, it is not always an inhabitant of swamps, and, second, I have never found it with the cap rimose. It may be added as a presumptive distinguishing feature that Fries makes no mention of the very noticeable and long persistent odor emitted by the drying and dried plants. The further fact that our plant has never yet been found with white milk, even in its upland growth, leads to the conclusion that it is certainly not a degenerate form but a species constant in its milk character, and in its decided and persistent odor and therefore worthy of specific distinction.

Entoloma grande Pk.

GRAND ENTOLOMA

PLATE II9, FIG. 1-5

Pileus fleshy, thin toward the margin, glabrous, convex becoming nearly plane, often broadly umbonate, sometimes rugosely wrinkled about the umbo, moist in wet weather, yellowish white or grayish brown, flesh white, odor and taste at first farinaceous, then sometimes leaving a disagreeable sensation in the mouth; lamellae broad, subdistant, slightly adnexed, whitish becoming pink; stem equal or nearly so, solid, slightly fibrous externally, mealy at the top, white; spores angular, .0003-.0004 of an inch in diameter (8-10 μ).

The grand entoloma is a large but rare mushroom. It has been found in a single locality near Albany twice in 13 years. It has been found once in the state of Vermont by Professor Burt. It is one of the few species of the genus *Entoloma* that have a farinaceous taste and odor. It grows in woods and occurs in August. It is

found single or in tufts. The cap is 2-6 inches broad, the stem 1.5-6 inches long and 3-12 lines thick. The cap is convex or somewhat bell shape, becoming nearly flat, whitish to grayish brown in color, its surface is smooth, and in large specimens it is sometimes umbonate and rugosely wrinkled about the umbo. Its flesh is white. The gills are at first whitish or grayish but as they mature they assume the pink color of the spores. They are rounded next the stem and but slightly attached to it. The stem is white or whitish, solid and often mealy at the top.

On account of the disagreeable sensation left in the mouth by tasting the uncooked cap it was thought that this mushroom would probably be found to be unwholesome. But actual experiment has shown that this character is destroyed by thorough cooking and that the mushroom is edible though less highly flavored than some others. Its scarcity makes it of but little importance.

Hebeloma album Pk.

WHITE HEBELOMA

PLATE II7, FIG. I-6

Pileus fleshy, thin, firm, convex becoming nearly plane or concave by the upcurving of the margin, glabrous, slightly viscid, white or yellowish white, flesh white, taste mild; lamellae thin, narrow, close, adnexed, whitish when young, becoming brownish ferruginous; stem equal, firm, solid, glabrous, slightly mealy at the top, white; spores subelliptic, .0005-.0006 of an inch long, .00025-.0003 broad (12-16 x 6-8 μ).

The white hebeloma is not a common mushroom but it is an excellent one for the table. It is gregarious in its mode of growth and occurs among fallen leaves and mosses or on naked damp soil in woods, and may be found in September and October. The cap is 1-2 inches broad, the stem 1-3 inches long and 2-3 lines thick. Generally the whole plant is white when young but the gills assume a brownish cinnamon or brownish rust color when mature. The edge of the gills is slightly excavated near the stem, to which they are narrowly attached. The cap is sometimes tinged with yellow and the stem is adorned at the top with white particles or a floccose mealiness. In State Museum Report 54, plate G, figures 1-7, the gills and spores are incorrectly colored. A new figure has therefore been prepared.

Boletus viridarius Frost

GREEN LAWN BOLETUS

PLATE 120, FIG. 1-10

Pileus fleshy, convex, viscid when moist, glabrous, dingy whitish, pale ochraceous reddish yellow or pale orange, inclining to reddish brown, flesh whitish or yellowish, unchangeable; tubes from plane to convex, usually slightly depressed around the stem, their mouths small or medium size, subrotund, the dissepiments at first whitish, becoming yellowish or yellowish brown when mature; stem equal or slightly tapering upward, solid, white or yellowish and distinctly reticulated above the slight white annulus, pallid, reddish or brownish below, whitish or yellowish within; spores oblong-fusiform, .0003-.0005 of an inch long, .00016-.00024 broad (8-12 x 4-6 μ).

Pileus 1-5 inches broad; stem 1-2.5 inches long, 3-6 lines thick.

Grassy ground near pine trees. September and October. Poughkeepsie. Miss H. L. Palliser.

This species is related to *Boletus flavus* With. by the stem being reticulate above the annulus, but it is far more variable in the color of the pileus and stem, and it also differs in the character of the margin of the pileus, which is often incurved and appendiculate by the remains of the white veil. In none of the specimens seen do I find any green hues, nor is anything said of green or greenish colors in the original description of the species by Mr Frost. We can therefore only infer that the specific name was suggested by the green grassy places in which this *Boletus* grows.

Its edible qualities have been tested both by Miss Palliser and myself and are considered excellent. The pileus is generally soiled by fragments of dirt or other matter, by reason of which it is better to remove the separable viscid cuticle before cooking. The plants vary in size. Those appearing in September are larger than those appearing in October. The tubes when young are whitish or pale yellow and where wounded assume a pale brownish or fawn color; when older they become brownish yellow and wounds assume a darker brown hue. The veil is white and in the later specimens appears to be more fully developed and more persistent than in the earlier ones. Its fragments in the later ones often adhere to the margin of the cap.

NEW SPECIES OF EXTRALIMITAL FUNGI

✓ *Amanita morrisii*

PLATE W, FIG. I-4

Pileus fleshy, subcampanulate becoming broadly convex, viscid when moist, glabrous, even on the margin, with a separable pellicle, dark grayish brown or blackish brown, becoming a little paler with age and with the escape of moisture, flesh white; lamellae thin, close, narrow, rounded behind, slightly adnexed, white; stem equal or slightly tapering upward, slightly bulbous at the base, solid or stuffed, slightly floccose, sometimes grayish and striate at the top, usually white, annulus double, radiately striate above, whitish buff beneath, the slight volva soon breaking into fragments and disappearing or occasionally partly adhering to the lower part of the stem; spores subglobose or broadly ellipsoid, $8-10 \times 6-8 \mu$.

Pileus 5-10 cm broad; stem 8-14 cm long, 12-20 mm thick.

In black vegetable mold among mosses. Natick swamp, Mass. September. G. E. Morris.

Pileus carnosus, subcampanulatus, deinde late convexus, viscidus, glaber, margine leve, pellicula separabile, griseo-brunneus seu atro-brunneus, in senectute vel quum siccus pallidior, carne alba; lamellae tenues, confertae, angustae, leviter adnexae, albae; stipes aequalis, vel sursum attenuatus, solidus farctusve, minute flocculosus, aliquando griseus et ad apicem striatus, vulgo albus, annulus crassus, mollis, supra radiate striatus et alba, infra luteolus, volva in fragmenta mox frangens et vanescens, seu stipitis parti inferiori in fragmentis rare adhaerens; sporae subglobosae vel ellipsoideae, $8-10 \times 6-8 \mu$.

✓ *Agaricus eludens*

PLATE X, FIG. 6-13

Pileus thin, ovate, broadly conic or subcampanulate, sometimes becoming broadly expanded, brown when young, becoming whitish and covered with brown fibrillose squamules, the center smooth, brown, the young margin surpassing the lamellae, flesh white changing to reddish where wounded; lamellae thin, close, narrow, free, whitish becoming bright pink, then chocolate brown and finally black or blackish brown; stem firm, nearly equal or sometimes thickened at the base, often slightly bulbous, fibrous, silky, white, stuffed with a hollow tube, internally white, changing to blood red

where wounded, then to brown or black, annulus thick, persistent, white; spores subglobose or ellipsoid, $5-7 \times 4-5 \mu$.

Pileus 2.5-10 cm broad; stem 2.5-7 cm long, 4-8 mm thick.

Cespitose or single, often in clusters of many individuals. On dumping ground near Trenton, N. J. September. E. B. Sterling.

The pileus closely resembles that of *Agaricus placomyces* Pk., but the mushroom differs in its commonly tufted mode of growth, the darker color of its mature pileus, the thicker and more persistent annulus, the distinct hollow tube of the stem and specially in the change of color of the wounded flesh and stem. It is also closely allied to *Agaricus approximans* Pk. from which it may be separated by its lamellae becoming pink before they assume the brown color of maturity. The discoverer pronounces it edible but says its flavor is less agreeable than that of *Agaricus campester* L.

Pileus tenuis, ovatus, late conicus seu subcampanulatus, aliquando late expansus, quum juvenis brunneus deinde albidus, squamulis brunneis fibrillosis tectus, centro glaber, brunneus, margine juvene lamellas excedente, carne alba, ubi vulnerata rufescente; lamellae tenues, confertae, angustae, liberae, albiae, mox incarnatae, deinde nigro-brunneae; stipes firmus, subaequalis, aliquando basi incrassatus saepe leviter bulbosus, fibrosus, sericeus, albus, tuba cava farctus, carne alba, ubi vulnerata sanguinea, deinde brunnea vel nigra, annulus crassus, persistens, albus; sporae subglobosae vel ellipsoideae, $5-7 \times 4-5 \mu$.

Russula blackfordae

PLATE Z, FIG. 9-13

Pileus fleshy but thin, broadly convex or nearly plane, viscid when moist, striate on the margin, whitish or pale gray, brown in the center, flesh white, taste mild; lamellae thin, narrow, close, adnate, pale yellow or cream color; stem equal, glabrous, stuffed or hollow, white; spores pale yellow, globose, $8-9 \mu$ broad.

Pileus about 2.5 cm broad; stem about 2.5 cm long, 4-6 mm thick.

Ellis, Mass. October. Mrs E. B. Blackford.

This species differs from *Russula fallax* (Schaeff.) Sacc. in the color of the pileus, the closer and yellowish lamellae, the mild taste and the color of the spores. The viscid pellicle of the pileus is separable. The species belongs to the section *Fragiles*, second subsection.

Pileus carnosulus, late convexus subplanusve, quum humidus, viscidus, margine striatus, albidus, pallido-griseusve, centro brunneus, carne alba, sapore miti; lamellae tenues, angustae, confertae, adnatae, flavidae vel cremeae; stipes aequalis, glaber, farctus vel cavus, albus; sporae globosae, flavae, 8-9 μ latae.

Russula serissima

Pileus fleshy, thin, fragile, convex becoming nearly plane or centrally depressed, viscid when moist, glabrous, with the margin even or sometimes obscurely striate when old, variable in color, pale olive-green or brownish purple, sometimes spotted in the center, occasionally pruinose, flesh white or whitish, taste mild or slightly and tardily acrid, odor in the dried or drying plant strong, unpleasant, persistent; lamellae thin, close, 4-8 mm broad, narrowed behind, adnexed sometimes seceding from the stem, cream color or buff, becoming dingy or smoky in drying; stem equal or tapering upward, solid but spongy within, white, both it and the flesh assuming a somewhat smoky hue in drying; spores subglobose, buff yellow, 10-12 x 8-10 μ .

Pileus 5-7 cm broad; stem 4-7 cm long, 8-20 mm thick.

Under fallen leaves in woods. Ellis, Mass. October. Mrs E. B. Blackford and G. E. Morris.

The *pileus* varies in color as does the *pileus* of *Russula variata* Banning and *Russula squalida* Pk. It is very close to the latter, from which it scarcely differs except in its viscid *pileus*, its late occurrence, its lamellae and flesh not changing color where wounded and specially in the color of the spore print.

Pileus carnosus, tenuis, fragilis, convexus, deinde subplanus vel centro depressus, quum humidus viscidus, glaber, margine levis aliquando in senectute leviter striatus, olivaceus vel brunneo-purpureus, aliquando centro maculatus, rare pruinosis, carne alba albidave, sapore miti vel leviter tardeque acri, odore ingrato, persistente; lamellae, tenues, confertae, 4-8 mm latae, adnexae, cremeae vel luteolae, deinde fumidae; stipes aequales vel sursum attenuatus, solidus, intra spongiosus, albus, deinde fumosus; sporae subglobosae, luteo-flavae, 10-12 x 8-10 μ .

Lactarius bryophilus

PLATE X, FIG. 1-5

Pileus thin, broadly convex or nearly plane, with an even margin, sometimes slightly umbonate, very viscid or glutinous, reddish

becoming subochraceous sometimes with one or two narrow orange zones near the margin, flesh white, taste mild, milk watery, scanty; lamellae unequal, close, adnate, whitish becoming ochraceous buff; stem soft, equal, glabrous, stuffed or hollow, colored like or a little paler than the pileus; spores subglobose, 6-8 μ in diameter.

Pileus 1-4 cm broad; stem 1.5-3.5 cm long, 4-8 mm thick.

Among mosses in swamps. Natick, Mass. September and October. G. E. Morris and S. Davis.

A very rare species hitherto known from no other locality, and only sparingly found in this one. It may be easily recognized by its small size, very viscid subochraceous pileus, mild taste and watery, unchangeable milk. In one or two cases very young specimens have appeared to have white milk, but in mature specimens the milk is constantly watery. This is doubtless its normal color.

Pileus tenuis, late convexus subplanusve, margine levis, subumbonatus, viscosus, rufus deinde subochraceus, aliquando juxta marginem zona angusta auratiaca ornatus, carne alba, sapore miti, lacte aquoso, parco; lamellae inaequales, confertae, adnatae, albidae, deinde flavo-ochraceae; stipes mollis, aequalis, glaber, farctus cavusve, colore pileo similis vel pallidior; sporae subglobosae, 6-8 μ latae.

Naucoria sphagnophila

Pileus thin, convex becoming nearly plane, minutely appressed tomentose and sometimes flocculose squamulose, hygrophanous, when young and moist tinged with flesh color, becoming buff white in drying, grayish ochraceous or rusty brown when mature; lamellae thin, narrow, subsinuate, close, unequal, uneven on the edge, yellowish becoming ferruginous; stem equal, flexuous, solid or at length hollow, yellowish with a slight floccose tomentum at the top, white tomentose at the base; spores ellipsoid, 8-9 x 4-5 μ .

Pileus 1.2-2.4 cm broad; stem 2.5-4.5 cm long, 2-3 mm thick.

Gregarious. In sphagnum in a swamp. Stow, Mass. July. S. Davis.

Pileus tenuis, convexus vel subplanus, minute tomentosulus, aliquando floccoso-squamulosus, hygrophanus, in juventate subincarnatus, in maturitate griseo-ochraceus vel ferrugineus; lamellae tenues, angustae, subsinuatae, confertae, inaequales, acie erosae, flavidae, deinde ferruginae; stipes aequalis, flexuosus, solidus, deinde fistulosus, luteolus, ad apicem minute floccoso-tomentosulus, basi albido-tomentosus; sporae ellipsoideae, 8-9 x 4-5 μ .

Cortinarius ferrugineo-griseus

PLATE Y, FIG. 1-4. PLATE Z, FIG. 1-3

Pileus convex or nearly plane, sometimes with the thin margin upcurved and then appearing centrally depressed, hygrophanous, brownish ferruginous when moist, gray or whitish gray when the moisture has escaped, flesh whitish; lamellae 4-6 lines broad, moderately close, adnexed, appearing free in the dried plant, pale cinnamon or clay color when young, brownish cinnamon when mature; stem equal, abruptly bulbous at the base, solid or stuffed, silky fibrillose, sometimes colored like but paler than the pileus, sometimes shining, variable in color, whitish below and violet tinted above or entirely violaceous, violaceous within; spores ellipsoid and commonly uninucleate, $10-12 \times 7-8 \mu$.

Pileus 3.5-10 cm broad; stem 3.5-8.5 cm long, 6-20 mm thick.

Under pine trees near Natick swamp, Mass. October. G. E. Morris.

The growing plant is often covered with pine needles. It belongs to subgenus *Hydrocybe* and is closely allied to *Cortinarius saturninus* Fr., from which it may be separated by its pileus fading to grayish white and by its solid stem often abruptly bulbous. It also differs in its habitat and in its larger spores.

Pileus convexus vel subplanus, aliquando centro depressus, margine recurvato, hygrophanus, quum humidus ferrugineo-brunneus, quum siccus albido-griseus griseusve, carne albida; lamellae subconfertae, adnexae, 8-12 mm latae, in juventute cinnamomeae vel argillaceae, in maturitate brunneo-cinnamomeae; stipes aequalis, basi abrupte bulbosus, sericeo-fibrillosus, nitidus, solidus, infra albidus, supra violaceus, vel omnino violaceus, aliquando in colore pileo similis sed pallidior, interne albidus vel violaceus; sporae ellipsoideae, vulgo uninucleatae, $10-12 \times 7-8 \mu$.

Cortinarius acutoides

PLATE Z, FIG. 4-8

Pileus submembranous, conic or subcampanulate, acute or acutely umbonate, hygrophanous, at first pale chestnut color floccose and white margined by the fibrils of the veil, after the escape of the moisture whitish and silky fibrillose; lamellae narrow, ascending, adnexed, subdistant, yellowish cinnamon; stem solid or with a small hollow, white, becoming whitish like the pileus; spores ellipsoid, $8-10 \times 6-7 \mu$.

Pileus 8-16 mm broad; stem 2.5-5 cm long, 2-3 mm thick.

Swamps. Ellis, Mass. October. Mrs E. B. Blackford.

Closely allied to *Cortinarius acutus* (Pers.) Fr., from which it differs in the darker color of the young moist pileus and whiter color of the mature dry pileus, the white color of the young stem, the adnexed lamellae, and specially by the larger spores and absence of striae from the moist pileus. This may be the plant mentioned in *Sylloge* as a variety of *Cortinarius acutus* (Pers.) Fr.

Pileus submembranaceus, conicus subcampanulatusve, acutus vel acute umbonatus, hygrophanus, primus pallido-castaneus, velo albo floccosus, margine albido, quum siccus albescens, sericeo-fibrillosus; lamellae angustae, ascendentes, adnexae, subdistantes, flavido-cinnamomeae; stipes solidus seu leviter cavus, albus deinde pileo in colore similis; sporae ellipsoideae, 8-10 x 6-7 μ .

✓ *Clavaria lavendula*

Tufts 2.5-4 cm high, densely and subdichotomously branched, the branches compressed, thin, lilac pink when moist, pruinose when dry, the ultimate ones often bidentate, axils rounded; spores minute 6-8 x 3-4 μ .

Chestnut grove. Stow, Mass. July. S. Davis.

This species is related to *Clavaria amethystina* Bull., but it differs in its flattened branches and smaller spores.

Caespites 2.5-4 cm alti, dense et subdichotome ramosissimi, rami tenues, numerosi, compressi, quum humidi lavenduli, sicci, pallidiores pruinosisque, ramuli ultimi saepe bidentati; sporae ellipsoideae, 6-8 x 3-4 μ .

✓ *Clavaria pallescens*

Clubs simple, loosely cespitose or gregarious, 2.5-4 cm tall, clavate, soft, fragile, obtuse, pale buff fading to whitish, sometimes minutely rugulose, stuffed or hollow, pale yellow within; stem short, glabrous, 2-4 mm long, pale yellow; spores oblong or elliptic, white, 9-12 x 6-8 μ .

Dry gravelly soil near *Kalmia angustifolia* L. South Acton, Mass. October. S. Davis and G. E. Morris.

This species is allied to *Clavaria ligula* Fr. from which it differs in its smaller size, in its color becoming whitish or paler with age or in drying, but being lemon-yellow and more persistent

within, in its glabrous lemon-yellow stem and in its broader spores. It is apparently a rare but very distinct species.

Clavae simplices, laxe caespitosae vel gregariae, 2.5-4 cm longae, clavatae, molles, fragiles, obtusae, luteolae, deinde albescentes, aliquando minute rugulosae, farctae cavaeve, intra flavae; stipes 2-4 mm longus, glaber, flavidus; sporae oblongae vel ellipsoideae, albae, 9-12 x 6-8 μ .

NEW YORK SPECIES OF INOCYBE

Inocybe Fr.

Veil universal, subfibrillose, concrete with the cuticle of the pileus, often free on the margin, webby; lamellae subsinuate (rarely adnate or decurrent) changing color, not cinnamon pulverulent; spores even, angular or rough, more or less brownish ferruginous. *Sylloge* 5:762

The species of this genus are generally of small or medium size. They were formerly included by Fries in the genus *Hebeloma*, from which the universal veil concrete with the commonly dry pileus specially distinguishes them. The prevailing color of the pileus is brown in some of its shades. In no other genus of the Agaricaceae is it more necessary to make use of the microscope in the identification of the species, for the external resemblance in some is so close that microscopic examination of the spores can not safely be omitted. The presence or absence of cystidia is also a character of some importance in the classification and identification of the species. Nearly all the species are terrestrial, some growing in woods, others in pastures and open places. A few occur on the ground and on decaying wood also. They have been distributed in five sections for convenience of study and the better understanding of their relations to each other. One author has instituted a genus depending on the rough spore character but it does not seem to find much favor among mycologists.

A microscopic examination of the spores would be necessary in such a case before even the generic identification could be made. Many of our species are rare or local, having been found but once and in a single locality.

In the following pages the arrangement of the sections as given in *Sylloge* has been followed. The following key to the sections is based on external characters and indicates the prominent characteristic of each section.

KEY TO THE SECTIONS

- Pileus and stem both squamose.....Squarrosae
 Pileus and stem not both squamose.....1
 1 Cuticle of the pileus lacerated or cracked.....2
 1 Cuticle of the pileus continuous.....3
 2 Pileus squamose or fibrillose lacerated.....Lacerae
 2 Pileus radiately rimose and fibrillose.....Rimosae
 3 Pileus not viscid.....Velutinae
 3 Pileus viscid.....Viscidae

Squarrosae

Pileus at first squamose or squarrosely squamose; stem squamose, colored like the pileus, both commonly some shade of brown.

This section differs from the others in having the pileus and stem alike in color and both squamose or squamulose.

KEY TO THE SPECIES

- Spores even.....1
 Spores not even.....4
 1 Pileus dark brown.....2
 1 Pileus not dark brown.....3
 2 Pileus 2.5-5 cm broad, scales persistent.....calamistrata
 2 Pileus 1.5-2.5 cm broad, scales subdeciduous.....mutata
 3 Pileus tawny, stem hollow, fibrillose squamulose.....fibrillosa
 3 Pileus subochraceous, stem solid, squamulose.....unicolor
 4 Pileus 2.5 cm broad.....stellatospora
 4 Pileus less than 2.5 cm broad.....lanuginosa

Inocybe calamistrata Fr.

CURVED SCALE INOCYBE

Sylloge 5: 762

Pileus fleshy, thin, campanulate or convex, obtuse, squarrosely squamose, brown or dark brown, flesh whitish, reddish where wounded; lamellae close, adnexed, whitish becoming ferruginous, the edge thick, whitish; stem equal, tough, solid, squarrosely squamose, brown, bluish at the base; spores oblong or ellipsoid, even. 10-15 x 5-6 μ .

Pileus 1-3 cm broad; stem 3-7 cm long, 2-4 mm thick.

Damp places under trees or bushes. Albany, Essex and Warren counties. August and September.

This species is well marked by the recurved scales of the pileus and stem and the bluish tint at the base of the stem. The European plant is described as having a strong odor but this character is scarcely noticeable in the American plant.

***Inocybe mutata* (Pk.) Mass.**

CHANGED INOCYBE

Ag. (*Hebeloma*) *mutatus* Pk. N.Y. State Mus. Rep't 24, p.69

Pileus thin, broadly conic or convex, obtuse or slightly and broadly umbonate, at first covered with erect or recurved scales which at length disappear except at the center, dark brown; lamellae broad, close, rounded at the stem, deeply sinuate, adnexed, ferruginous brown, crenulate on the edge; stem slender, equal, solid, floccosely scaly, often curved at the base, brown; spores ellipsoid, even, $9-11 \times 5-6$.

Pileus 1.5-2.5 cm broad; stem 5-7 cm long, about 2 mm thick.

Damp ground in woods. Ulster co. July.

The species is closely related to *Inocybe calamistrata* Fr. from which it may be separated by its smaller size, scales disappearing from the margin, absence of bluish tints from the base of the stem and shorter spores. The changed appearance of the pileus caused by the vanishing scales of the margin is suggestive of the specific name.

***Inocybe fibrillosa* Pk.**

FIBRILLOSE INOCYBE

N. Y. State Mus. Rep't 41, p.65

Pileus thin, convex or nearly plane, obtuse or subumbonate, densely fibrillose, tawny, generally a little darker in the center and there adorned with appressed fibrillose scales; lamellae close, adnate, yellowish or yellowish olivaceous becoming brownish cinnamon; stem equal, hollow, fibrillosely squamose, colored like or a little paler than the pileus; spores ellipsoid, even, $8-10 \times 5-6 \mu$.

Pileus 2-3.5 cm broad; stem about 2.5 cm long, 2-4 mm thick.

Damp mossy banks in woods. Albany co. August. A rare species.

***Inocybe unicolor* Pk.**

ONE COLORED INOCYBE

N. Y. State Mus. Rep't, 50, p.104

Pileus conic or very convex becoming broadly convex or nearly plane, tomentosely squamulose, pale ochraceous or grayish ochraceous, flesh white; lamellae broad, subdistant, subventricose, pale ochraceous becoming tawny brown; stem slender, firm, equal, flexu-

ous, solid, squamulose, colored like the pileus; spores ellipsoid, even, $8-12 \times 5-6 \mu$.

Pileus 2-2.5 cm broad; stem 2.5-3 cm long, 2-4 mm thick.

Clay soil. Albany co. July. Rare.

This species approaches *Inocybe subochracea* (Pk.) Mass. in color, but it differs in having the stem squamulose and colored like the pileus and in its larger spores.

Inocybe stellatospora (Pk.) Mass.

STELLATE SPORE INOCYBE

Ag. (*Hebeloma*) *stellatosporus* Pk. N. Y. State Mus. Rep't 26, p.57

Pileus thin, convex or nearly plane; dry, covered with erect or curved scales, dark brown; lamellae close, adnate, pallid becoming brown or slightly rusty brown; stem equal, firm, solid, squamose, colored like the pileus; spores subglobose, nodulose, $7-8 \mu$ in diameter, cystidia $70-80 \times 14-20 \mu$.

Pileus about 2.5 cm broad; stem 4-5 cm long, about 2 mm thick.

In woods. Lewis co. September.

In size and color this species resembles *Inocybe mutata* (Pk.) Mass. but it is easily distinguished by its persistent scales on the pileus and by its nodulose spores.

Inocybe lanuginosa (Bull.) Karst.

WOOLLY INOCYBE

Ag. (*Inocybe*) *nodulosporus* Pk. N. Y. State Mus. Rep't 32, p.28

Pileus thin, hemispheric or convex, obtuse, floccosely squamose, cervine brown or umber color, the scales of the disk usually erect; lamellae close, broad, ventricose, rounded at the stem, pallid becoming ferruginous cinnamon, white and crenulate on the edge; stem slender, equal, solid, flexuous, tomentosely squamulose, colored like the pileus; spores globose or subellipsoid, nodulose, $6-8 \mu$ in diameter or $8-10 \times 8 \mu$, cystidia ellipsoid, $30-40 \times 16-20 \mu$.

Pileus 1-2 cm broad; stem 2-2.5 cm long, 2 mm thick.

Decaying wood in woods. Saratoga co. August.

European authors do not all agree concerning the character of the spores of this species, describing them as even, angular and acutely warty. In our specimens, which were at first supposed to be distinct, they are as here described. In other characters the

agreement with the description of *Inocybe lanuginosa* (Bull.) Karst. as given in *Sylloge* is so close that it seems best to refer our plant to this species.

Lacerae

Cuticle of the pileus squamose or fibrillose lacerated; stem paler than the pileus.

KEY TO THE SPECIES

- Spores even.....1
- Spores angular or nodulose.....2
- 1 Pileus brown, fibrillose squamulose.....infelix
- 1 Pileus ochraceous buff, rimose squamose in the center....squamosodisca
 - 2 Spores angular.....maritimoides
 - 2 Spores nodulose.....3
- 3 Pileus brown or grayish brown.....diminuta
- 3 Pileus tawny or ochraceous.....subfulva

Inocybe infelix Pk.

UNFORTUNATE INOCYBE

Ag. (*Inocybe*) *infelix* Pk. N. Y. State Mus. Rep't 32, p.29

Pileus campanulate broadly convex or nearly plane, subumbonate, floccosely squamulose, grayish brown or umber, flesh whitish; lamellae close, adnexed, ventricose, broad, whitish becoming brownish ferruginous; stem equal, solid, silky fibrillose, pallid or whitish above, generally brownish toward the base, pruinose at the top; spores oblong, even, 10-15 x 5-6 μ , cystidia flask shape, 40-60 x 15-20 μ .

Pileus 1.5-2.5 cm broad; stem 2-5 cm long, 2-4 mm thick.

Naked sterile soil or among mosses. Albany, Saratoga, Essex and Hamilton counties. May to August.

Var. **brevipes** Pk. N. Y. State Mus. Bul. 2, p.13.

Pileus scarcely exceeding 1.5 cm broad, stem about 1.5 cm long.

This is a common and variable species, but it is easily recognized by its persistently squamulose brown pileus and its oblong even spores. In wet weather the cuticle of the pileus is often more lacerated than in dry weather. The umbo is sometimes wanting. The plants occur throughout the season when the weather conditions are favorable. It is gregarious in its mode of growth.

Inocybe squamosodisca Pk.

SCALY DISK INOCYBE

N. Y. State Mus. Bul. 75, p.18

Pileus fleshy, firm, convex, fibrillose on the margin, rimosely squamose in the center, ochraceous or ochraceous buff, flesh whitish or yellowish white; lamellae broad, moderately close, adnate, pale ochraceous becoming darker with age; stem equal, solid, fibrillose, colored like or a little paler than the pileus; spores ellipsoid, even, $8-10 \times 5-6 \mu$.

Pileus 2.5-5 cm broad; stem about 2.5 cm long, 4-6 mm thick.

Gregarious. Under pine trees. Hamilton co. August. Rare.

The scales of the pileus are caused by the cracking of the cuticle.

Inocybe maritimoides Pk.

MARITIMOID INOCYBE

N. Y. State Mus. Rep't 38, p.87

Pileus subconic or convex, obtuse or slightly umbonate, densely squamulose in the center, fibrillose on the margin, dark brown; lamellae close, adnexed, ventricose, whitish becoming brownish ochraceous; stem equal, solid, fibrillose, colored like but paler than the pileus; spores irregular, angular, ovate or ellipsoid, $7-9 \times 5-6 \mu$, cystidia $40-55 \times 12-20 \mu$.

Pileus 1.5-2.5 cm broad; stem about 2.5 cm long, 4 mm thick.

Sandy soil in woods. Albany co. October. Rare.

It resembles *Inocybe maritima* Fr. but is separated from it by the pileus which is not hygrophane and by the spores which are smaller and angular but not nodulose.

Inocybe diminuta Pk.

SMALL INOCYBE

N. Y. State Mus. Bul. 105, p.23

Pileus thin, hemispheric becoming convex or nearly plane, squamose with erect or squarrose hairy scales in the center, fibrillose on the margin, grayish brown; lamellae subdistant, broadly sinuate, adnexed, ventricose, whitish becoming rusty brown; stem short, firm, solid, silky fibrillose, whitish above, grayish brown and slightly squamulose toward the base; spores subglobose, nodulose, $8-10 \times 6-8 \mu$, cystidia $40-50 \times 12-20 \mu$.

Pileus 6-12 mm broad; stem 8-16 mm long, 2 mm thick.

Bare compact soil in roads in woods. Suffolk co. August. Rare.

It appears like a dwarf form of *Inocybe infelix* Pk. but it is very distinct in the character of the spores.

Inocybe subfulva Pk.

TAWNY INOCYBE

N. Y. State Mus. Rep't 41, p.66

Pileus broadly conic or subcampanulate, becoming convex or nearly plane, subumbonate, fibrillose squamose, tawny ochraceous; lamellae broad, close, adnexed, ventricose, pallid becoming tawny cinnamon; stem equal, firm, solid, fibrous striate, obscurely pruinose, colored like but paler than the pileus; spores globose or ellipsoid, stellately nodulose, 8-10 μ in diameter or 10-12 x 7-8 μ , cystidia 40-80 x 10-15 μ .

Pileus 1.5-3 cm broad; stem 2.5-5 cm long, 2-4 mm thick.

Sandy soil in fields. Albany co. August.

This species is closely allied to *Inocybe gaillardii* Gill, from which it may be distinguished by its larger size, solid stem and variable spores. The scales of the center of the pileus are often erect but not squarrose.

Rimosae

Pileus radiately fibrillose, soon radiately rimose, sometimes adorned with appressed scales; stem fibrillose, white or whitish or slightly tinged with the color of the pileus.

The species of this section are easily recognized by the radiately cracking of the cuticle of the pileus and the pale color of the stem.

KEY TO THE SPECIES

- * Spores even.....1
- Spores angular or slightly nodulose.....7
- Spores distinctly nodulose.....8
- 1 Pileus umbonate.....2
- 1 Pileus not umbonate.....5
 - 2 Cuticle peeling in scales or patches.....excoriata
 - 2 Cuticle not peeling.....3
- 3 Pileus pale brown.....pallidipes
- 3 Pileus yellowish brown.....rimosa
- 3 Pileus some other color.....4
 - 4 Pileus fawn color.....eutheloides
 - 4 Pileus grayish fawn or chestnut.....eutheloides
- 5 Pileus yellowish brown.....rimosa

- 5 Pileus gray or grayish.....6
 6 Young lamellae whitish.....griseoscabrosa
 6 Young lamellae pale violaceous.....violaceifolia
 7 Pileus chestnut color.....castanea
 7 Pileus brown or dark brown.....umboninota
 7 Pileus brownish with a whitish center.....albodisca
 8 Pileus tawny gray.....rigidipes
 8 Pileus brownish.....asterospora

***Inocybe excoriata* Pk.**

EXCORIATE INOCYBE

N. Y. State Mus. Bul. 75, p.16, pl.O, fig.14-19

Pileus fleshy, broadly conic becoming broadly convex, umbonate, fibrillose or fibrillosely squamulose, slightly silky or tomentose on the margin, grayish brown, the cuticle often cracking and peeling, flesh white; lamellae close, narrow, adnexed, with a decurrent tooth, white becoming brownish gray, whitish and crenulate on the edge; stem equal, solid, silky fibrillose, white or whitish; spores ellipsoid, even, 8-10 x 5-6 μ , cystidia flask shape, 50-60 x 12-20 μ .

Pileus 2.5-5 cm broad; stem 2.5-5 cm long, 4-6 mm thick.

Among fallen leaves in woods. Hamilton co. August.

The surface of the pileus cracks radiately and thereby indicates the section to which the species belongs. A slight whitish webby veil is present in the young plant.

***Inocybe pallidipes* E. & E.**

PALE STEM INOCYBE

Jour. Myc. 5, p.24

Pileus conic or campanulate becoming expanded, umbonate, fibrillosely squamose, innately or subrimosely scaly on the disk, subrimose on the margin, brown or pale brown; lamellae subclose, rather broad, ascending, becoming ventricose, adnate with a decurrent tooth, pallid becoming clay color or watery cinnamon; stem solid, slightly narrowed and mealy above, loosely fibrillose below, subbulbous, white, white tomentose at the base; spores unequally ellipsoid, even, 7-10 x 5-6 μ , cystidia ventricosely fusoid or flask shape, 40-60 x 14-20 μ .

Pileus 2-3 cm broad; stem 2.5-5 cm long, 2-4 mm thick.

Decaying wood and vegetable mold. Warren co. July.

This species may be recognized by its umbonate pileus and persistently white stem. The umbo is sometimes more highly colored

than the rest of the pileus. From *Inocybe eutheloides* Pk. it may be distinguished by its lamellae and white stem.

***Inocybe rimosa* (Bull.) Fr.**

CRACKED INOCYBE

Sylloge 5:775

Pileus fleshy, thin, broadly conic or campanulate becoming expanded, obtuse or umbonate, silky fibrous, radiately cracking on the surface, yellowish brown; lamellae subclose, adnexed or nearly free, whitish becoming tan color or subferruginous; stem equal, firm, solid, mealy at the top, nearly glabrous, subbulbous, whitish; spores ellipsoid, even, $10-12 \times 5-6 \mu$, cystidia very rare, $60-65 \times 15-18 \mu$.

Pileus 2.5-5 cm broad; stem 2-5 cm long, 4-6 mm thick.

Ground in woods and open places. Albany, Franklin and Ulster counties. August and September.

This is a very variable species but one which is generally recognizable by the radiately cracking of the surface of the pileus.

Var. **parva** Pk. Very small; the pileus rarely more than 2 cm broad, the cuticle obscurely cracking or sometimes continuous.

Var. **cuspidata** Pk. Pileus with a very prominent narrow subacute or cusplike umbo.

***Inocybe eutheles* (B. & Br.) Sacc.**

MAMMILLATE INOCYBE

Sylloge 5:776

Pileus thin, campanulate becoming expanded, distinctly umbonate, silky, shining, subsquamulose, pale fawn color; lamellae rather narrow, adnate, pallid becoming subferruginous, whitish crenulate on the edge; stem slender, solid, equal, fibrous, pallid or whitish; spores ellipsoid, even, $10-15 \times 6-8 \mu$, cystidia very rare, $60-65 \times 15-20 \mu$.

Pileus 2.5-4 cm broad; stem 3-6 cm long, 2-3 mm thick.

Ground. Albany co.

This species has been found but once and is apparently very rare. The European plant is said to have a farinaceous odor and spores 7-10 μ long. In our plant the spores are longer and the odor was not noticed, but in other respects the agreement is fairly good.

Inocybe eutheloides Pk.

EUTHELOID INOCYBE

N. Y. State Mus. Rep't 32, p.29

Pileus thin, campanulate becoming expanded, distinctly umbonate, silky fibrillose, subrimose, varying from grayish cervine to chestnut color, sometimes squamulose on the disk; lamellae close, rather broad, ventricose, narrowed toward the stem, adnexed, whitish becoming brownish ferruginous, white and crenulate on the edge; stem equal, subflexuous, solid, fibrillose, pallid or whitish; spores unequally ellipsoid, uninucleate, even, $8-12 \times 5-6 \mu$, cystidia ventricose, $45-55 \times 12-16 \mu$.

Pileus 1.5-2.5 cm broad; stem 2.5-5 cm long, 2-4 mm thick.

Ground in woods. Onondaga co. September.

This differs from *Inocybe eutheles* (B. & Br.) Sacc. in its smaller size, darker color, adnexed lamellae, and slightly smaller spores.

Inocybe griseoscabrosa (Pk.) Mass.

GRAYISH INOCYBE

Ag. (*Hebeloma*) *griseoscabrosus* Pk. N. Y. State Mus. Rep't 26, p.57

Pileus hemispheric or convex, fibrillosely squamulose, cinereous, with margin whitish when young; lamellae broad, close, whitish becoming brownish ferruginous; stem firm, equal or slightly tapering downward, solid, fibrillose or slightly squamulose, whitish or grayish; spores ellipsoid, even, $9-12 \times 5-8 \mu$, cystidia ventricose, $40-60 \times 15-20 \mu$.

Pileus 1-2 cm broad; stem 3-5 cm long, 2-3 mm thick.

Ground in woods. Albany co. October. Rare.

This small species has been found but once. It is peculiar in having a grayish or cinereous pileus with a white margin.

Inocybe violaceifolia Pk.

VIOLACEOUS GILL INOCYBE

N. Y. State Mus. Rep't 26, p.57

Pileus thin, convex or nearly plane, floccosely fibrillose, sub-squamulose, grayish buff; lamellae close, adnexed, pale violaceous becoming pale cinnamon; stem firm, solid, slender, fibrillose, white or whitish; spores ellipsoid, even, $8-10 \times 5-6 \mu$, cystidia $40-60 \times 15-20 \mu$.

Pileus 1-2 cm broad; stem about 2.5 cm long, 2mm thick.

Mossy ground in woods. Albany co. Rare.

This is a small pale species remarkable for the violaceous tint of the young lamellae. Sometimes the pileus is slightly umbonate. A webby veil is present in the young plant.

***Inocybe castanea* Pk.**

CHESTNUT INOCYBE

N. Y. State Mus. Bul. 75, p.16, Pl.O, fig.1-8

Pileus conic or convex, umbonate, radiately rimose, fibrillose, chestnut color; lamellae thin, narrow, close, adnate, whitish becoming brownish ferruginous; stem equal, hollow, subglabrous, pruinose or mealy at the top, often with a whitish tomentum at the base, colored like but paler than the pileus; spores subglobose, angular or slightly nodulose, 6-8 μ in diameter or 8 x 6 μ , cystidia subfusiform, 50-80 x 12-18 μ .

Pileus 1-2 cm broad; stem 2.3-5 cm long, about 2 mm thick.

Mossy ground under spruce and balsam fir trees. Hamilton co. August. Rare.

This is a well marked species and not easily confused with any other.

***Inocybe umboninota* Pk.**

UMBO MARKED INOCYBE

Ag. (*Inocybe*) *umboninotus* Pk. N. Y. State Mus. Rep't 38, p.87 in part

Pileus broadly campanulate becoming expanded, umbonate, fibrillose, slightly rimose, dark brown, the umbo sometimes darker than the rest of the pileus; lamellae close, adnate, sometimes slightly sinuate, whitish becoming brownish ferruginous; stem equal or slightly thickened at the base, solid, colored like but paler than the pileus, pruinose at the top; spores subglobose or ellipsoid, very slightly nodulose, 6-8 x 4-6 μ , cystidia 50-60 x 12-20 μ .

Pileus 1.5-2 cm broad; stem 2.5-5 cm long, 2-4 mm thick.

Ground in woods. Ulster co. September.

This species is closely related to *Inocybe asterospora* Quel. from which it may be separated by its less distinctly rimose pileus and by its smaller less globose and but slightly nodulose spores.

Inocybe albodisca Pk.

WHITE DISK INOCYBE

N. Y. State Mus. Rep't 51, p.290

Pileus conic or campanulate, umbonate, glabrous, whitish in the center when moist, elsewhere yellowish brown or lilac brown, paler when dry, slightly silky fibrillose, radiately rimose; lamellae subclose, adnexed, whitish becoming subferruginous; stem equal, solid, striate, slightly mealy or pruinose at the top, pallid; spores subglobose or ellipsoid, slightly nodulose, 6-8 μ in diameter or 8 x 6 μ , cystidia 40-60 x 14-20 μ .

Pileus 1.5-2.5 cm broad; stem 2-5 cm long, 3-4 mm thick.

Under spruce and balsam fir trees. Essex co. August.

The species is well marked by the whitish umbo or center of the pileus.

Inocybe rigidipes Pk.

RIGID STEM INOCYBE

N. Y. State Mus. Rep't 51, p.289

Pileus thin, convex or subcampanulate becoming convex, umbonate, squamulose, striate and slightly rimose on the margin when dry, tawny gray; lamellae broad, subdistant, narrowed toward the stem, slightly adnexed, tawny ochraceous; stem slender, flexuous, rigid, firm, solid, slightly pruinose, colored like the pileus; spores globose, strongly nodulose, 12 μ in diameter, cystidia 45-60 x 12-16 μ .

Pileus 1.5-2.5 cm broad; stem 3.5-6 cm long, about 2 mm thick.

Damp clay soil in shaded places. Albany co. August. Rare.

This species resembles *Inocybe subfulva* Pk. from which it may be distinguished by its globose spores and tawny gray pileus. It is also related to *Inocybe calospora* Quel. from which it differs in its tawny gray color, slightly adnexed lamellae, solid flexuous stem and larger spores.

Inocybe asterospora Quel.

STAR SPORE INOCYBE

Sylloge 5:780

Pileus campanulate becoming expanded, umbonate, radiately rimose, fibrillose, brown or brownish, the umbo often darker than the rest; lamellae close, dull cinnamon; stem equal, subbulbous, sub-

glabrous, solid, whitish or tinged with the color of the pileus; spores subglobose, nodulose, 8-12 μ in diameter, cystidia 40-70 x 12-20 μ .

Pileus 2-3 cm broad; stem 3-6 cm long, 2-3 mm thick.

Ground in woods. Fulton and Rensselaer counties. June and July.

It bears some resemblance to *Inocybe rimosa* (Bull.) Fr. but from that species it is at once separated by its nodulose subglobose spores. A form with brown cap and prominent umbo was formerly confused with *Inocybe umboninota* Pk. from which it is also best separated by its spores.

Velutinae

Cuticle of the pileus not cracking, covered with interwoven fibrils, becoming smooth or appressedly squamose, disk even; stem polished, smooth, whitish, mealy at the top.

KEY TO THE SPECIES

- | | |
|--|--------------|
| Spores even..... | 1 |
| Spores slightly nodulose..... | 5 |
| Spores distinctly nodulose..... | 7 |
| 1 Pileus white or whitish, rarely lilac tinted..... | 2 |
| 1 Pileus some other color..... | 3 |
| 2 Pileus 2-8 mm broad..... | comatella |
| 2 Pileus 1-2.5 cm broad..... | geophylla |
| 2 Pileus 2.5-7 cm broad..... | serotina |
| 3 Pileus pale ochraceous..... | subochracea |
| 3 Pileus pale tawny or brownish tawny..... | 4 |
| 4 Pileus pale tawny, umbonate..... | agglutinata |
| 4 Pileus brownish tawny, obtuse..... | subtomentosa |
| 5 Pileus 2.5-5 cm broad..... | fallax |
| 5 Pileus less than 2.5 cm broad..... | 6 |
| 6 Pileus whitish or pallid 1-2 cm broad..... | paludinella |
| 6 Pileus chestnut or subochraceous, 6-10 mm broad..... | subexilis |
| 7 Pileus blackish brown with a grayish margin when moist,
cinereous when dry..... | nigridisca |
| 7 Pileus whitish, often with a reddish brown umbo..... | infida |

Inocybe comatella Pk.

HAIRY CAP INOCYBE

N. Y. State Mus. Rep't 38, p.87, pl.2, fig.5-8

Pileus very thin, convex or expanded, clothed with whitish or grayish hairs, fimbriate on the margin; lamellae subdistant, adnexed, cinnamon; stem equal, solid, flexuous, pallid or reddish brown, slightly mealy at the top, slightly flocculose pruinose, with a

whitish mycelial tomentum at the base; spores subellipsoid, even, $8-10 \times 5-6 \mu$, cystidia $45-55 \times 12-20 \mu$

Pileus 4-8 mm broad; stem 1.5-2.5 cm long, about 1 mm thick.

Decaying wood and bark buried under fallen leaves. Fulton co. July. Rare. Found but once.

This is a very small but distinct species remarkable for the hairy covering of the pileus. The hairs on the margin are longer and coarser than the others. The habitat is unusual for an *Inocybe*.

***Inocybe geophylla* (Sow.) Fr.**

EARTHY LEAF INOCYBE

Sylloge 5:784

Pileus fleshy but thin, conic or ovate becoming expanded, umbonate, silky fibrillose, even, commonly white or whitish, rarely lilac; lamellae close, rather broad, ventricose, adnexed, white becoming clay color; stem equal, firm, stuffed, white, mealy at the top, spores ellipsoid, $8-10 \times 4-6 \mu$, cystidia $40-60 \times 12-20 \mu$.

Pileus 1.5-2.5 cm broad; stem 2.5-6 cm long, 2-4 mm thick.

Ground in woods. Common. August to October.

A fine but small species found mostly in woods and varying some in the color of the pileus. Var. *lilacinus* Pk. with lilac colored pileus fading to whitish when old appears to include both *Agaricus affinis* Pers. and *Agaricus geophilus* Pers.

***Inocybe serotina* Pk.**

LATE INOCYBE

N. Y. State Mus. Bul. 75, p.17

Pileus fleshy, firm, campanulate or convex becoming nearly plane, fibrillose on the margin, white or yellowish, flesh white; lamellae close, rounded at the stem, slightly adnexed, subventricose, whitish becoming brownish cinnamon; stem subequal, bulbous or sometimes narrowed at the base, solid, fibrous, white; spores oblong or ellipsoid, uninucleate, even, $12-16 \times 6-8 \mu$.

Pileus 2.5-6 cm broad; stem 2.5-6 cm long, 6-12 mm thick.

Sandy soil of Lake Ontario. Wayne co. October. Rare or local. Found but once.

This is one of the largest of our species. Its late appearance in the season is suggestive of the specific name. It is reported by its discoverer, E. B. Burbank, to be edible.

Inocybe subochracea (Pk.) Mass.

OCHRACEOUS INOCYBE

Ag. (Hebeloma) subochraceus Pk. N. Y. State Cab. Rep't 23, p.95

Pileus thin, conic or convex becoming expanded, commonly umbonate, fibrillose squamulose, ochraceous yellow; lamellae close, sinuate next the stem, adnexed, whitish becoming ferruginous or brownish ferruginous; stem equal, solid, slightly fibrillose whitish; spores ellipsoid, even, $8-10 \times 5-6 \mu$, cystidia $40-60 \times 12-16 \mu$.

Pileus 2-4 cm broad; stem 1-6 cm long, 2-4 mm thick.

Ground in woods and open places. August to October. Common.

Var. **burtii** Pk. N. Y. State Mus. Rep't 54, p. 167, pl. H, fig. 23-29. Veil distinct, webby, adhering to the margin of the pileus and to the stem, stem long, 5-7.5 cm, fibrillose; mature lamellae darker colored.

Inocybe agglutinata Pk.

AGGLUTINATE INOCYBE

N. Y. State Mus. Rep't 41, p.67

Pileus conic or campanulate becoming convex, umbonate, appressedly fibrillose, sometimes streaked or spotted by the colored fibrils, pale tawny, the umbo very prominent, brown; lamellae close, broad, ventricose, adnexed, whitish becoming brownish cinnamon, usually whitish on the edge; stem firm, solid, pruinose at the top, white or whitish above, tawny or brown toward the base, fibrillose; spores subovate or ellipsoid, even, $10-12 \times 5-6 \mu$, cystidia $40-60 \times 16-24 \mu$.

Pileus 1-2.5 cm broad; stem 2.5-5 cm long, 2-4 mm thick.

Under pine trees. Ulster co. September. Rare.

The fibrils of the pileus appear as if glued to its surface but the pileus is not viscid. The species is very similar to *Inocybe whitei* B. & Br. in general appearance, but it differs from that species in having a very prominent umbo, in the absence of viscosity from the pileus and in its larger spores.

Inocybe subtomentosa Pk.

SUBTOMENLOSE INOCYBE

N. Y. State Mus. Rep't 48, p.11

Pileus thin, dry, convex or plane, minutely tomentose, brownish tawny; lamellae thin, close, adnate, slightly sinuate, brownish tawny;

stem short, solid, slightly silky fibrillose, colored like or a little paler than the pileus, often with a conspicuous white mycelioid tomentum at the base; spores subellipsoid, even, $8-10 \times 5-7 \mu$.

Pileus 1.5-2.5 cm broad; stem 1.5-2.5 cm long, 2 mm thick.

Gravelly soil among fallen leaves. Clinton co. September. Rare.

This species is related to *Inocybe tomentosa* E. & E. from which it may be separated by the darker color of the pileus, the absence of an umbo and the larger spores. Its distinguishing features are its small size and minutely tomentose pileus of a uniform brownish tawny color.

Inocybe fallax Pk.

FALLACIOUS INOCYBE

N. Y. State Mus. Bul. 75, p.17, pl.O, fig.20-24

Pileus thin, campanulate or convex, umbonate, obscurely fibrillose, sometimes minutely and obscurely squamulose, whitish or whitish buff, subshining, the margin decurved and often splitting; lamellae thin, close, adnexed, pallid becoming brownish ferruginous; stem equal, flexuous, hollow, minutely mealy or pruinose, whitish; spores angular or slightly nodulose, $8-10 \times 6-8 \mu$, cystidia $40-50 \times 15-18 \mu$.

Pileus 2.5-5 cm broad; stem 5-7.5 cm long, 4-8 mm thick.

Among fallen leaves in woods. Hamilton co. August. Rare.

This species resembles large forms of *Inocybe geophylla* (Sow.) Fr. from which it may be separated by its spores.

Inocybe paludinella Pk.

MARSH INOCYBE

Ag. (*Inocybe*) *paludinellus* Pk. N. Y. State Mus. Rep't 31, p.34

Pileus thin, slightly convex, soon plane, umbonate, subfibrillose, whitish or pallid; lamellae narrow, close, adnate, whitish becoming subferruginous; stem slender, equal, colored like the pileus with a mass of white mycelium at the base; spores subangular, very slightly nodulose, $6-8 \times 5-6 \mu$, cystidia $45-60 \times 12-16 \mu$.

Pileus 1-2 cm broad; stem 2.5-5 cm long, 1-1.5 mm thick.

Gregarious. Low ground and wet places under bushes. Rensselaer co. August. Rare.

This species resembles small forms of *Inocybe geophylla* (Sow.) Fr. in color, but it is at once separated from that species by its slightly nodulose spores. From *Inocybe trechispora*

Berk. it differs in its smaller size, more slender stem, and in its pileus being neither viscid when moist nor silky when dry. Its lamellae are adnate and fairly bristle with numerous cystidia.

Inocybe subexilis Pk.

FEEBLE INOCYBE

Ag. (*Inocybe*) *subexilis* Pk. N. Y. State Mus. Rep't 38, p.87

Pileus thin, convex or subcampanulate becoming expanded, umbonate, fibrillose on the margin, pale chestnut becoming yellowish or subochraceous; lamellae narrow, close, adnexed, whitish becoming dingy ochraceous; stem equal, slender, flexuous, slightly striate, solid, minutely pruinose, pinkish becoming yellowish; spores subglobose, slightly nodulose, 5-6 μ in diameter, cystidia 45-60 x 12-15 μ .

Pileus 6-10 mm broad; stem 2-2.5 cm long, about 1 mm thick.

Damp mossy ground in woods. Fulton co. July. Rare.

Inocybe nigridisca Pk.

BLACK DISK INOCYBE

N. Y. State Mus. Rep't 41, p.67

Pileus thin, convex becoming nearly plane or centrally depressed, umbonate, minutely fibrillose and blackish brown with a grayish margin when moist, cinereous when dry; lamellae close, rounded at the stem, free or slightly adnexed, grayish becoming brownish ferruginous; stem slender, flexuous, firm, solid, minutely villose pruinose, reddish brown; spores globose or ellipsoid, nodulose, 5-6 μ in diameter or 7-8 x 5-6 μ , cystidia 45-50 x 12-15 μ .

Pileus 8-16 mm broad; stem 2.5-4 cm long, about 1 mm thick.

Damp places under ferns. Oswego co. June. Rare.

Its distinguishing features are its blackish brown pileus with grayish margin when moist, fading to cinereous when dry.

Inocybe infida (Pk.) Mass.

UNTRUSTY INOCYBE

Ag. (*Inocybe*) *infidus* Pk. N. Y. State Mus. Rep't 27, p.95

Pileus firm, campanulate or expanded, subumbonate, slightly squamulose on the disk, often split on the margin, whitish with umbo or disk often reddish brown; lamellae close, narrow, adnexed, pallid, becoming subcinnamon; stem equal or a little enlarged at

the base, furfuraceous at the top, hollow, white; spores subglobose, nodulose, $8-10 \times 6-8 \mu$, cystidia $40-60 \times 12-20 \mu$.

Pileus 1.5-2.5 cm broad; stem 3-5 cm long, 2-4 mm thick.

Mossy ground in low woods. Essex co. September.

The resemblance of this species to some forms of *Inocybe geophylla* (Sow.) Fr. is so close that it is important to have a knowledge of its spore characters in order to make a satisfactory determination. The specific name is suggestive of this fact. Sometimes the margin is so abundantly and deeply split that the radiating lobes give a stellate appearance to the pileus. *Inocybe commixta* Bres., *Inocybe umbratica* Quel. and *Inocybe leucocephala* Boud. are given as synonyms of this species by Massee.

This species has been reported as having caused a slight temporary illness in some members of a family who had specimens of it prepared for the table and partook of them. It is well therefore to consider it a poisonous or at least an unwholesome species.

Viscidae

Pileus viscid becoming smooth.

This section connects the genus *Inocybe* with the genus *Hebeloma*, the viscid pileus being common to it and *Hebeloma*. The character, "becoming smooth," does not rigidly apply in all cases, for in some of the species the pileus is more or less persistently silky or fibrillose or hairy on the margin.

KEY TO THE SPECIES

- Spores globose or subglobose.....*tricholoma*
- Spores not globose.....1
- 1 Spores nodulose*trechispora*
- 1 Spores not nodulose.....2
- 2 Pileus blackish brown in the center.....*fuscodisca*
- 2 Pileus not blackish brown in the center.....*vatricosoides*

Inocybe tricholoma (A. & S.) Fr.

HAIRY MARGIN INOCYBE

Paxillus strigosus Pk. N. Y. State Mus. Rep't 26, p.63

Pileus thin, broadly convex becoming nearly plane or slightly depressed in the center, subviscid, slightly hairy, specially on the subciliate margin, whitish; lamellae close, narrow, decurrent, whitish becoming brownish or subferruginous; stem equal, stuffed or solid

pruinose, whitish; spores subglobose, even or minutely nodulose, 4-5 μ in diameter.

Pileus 2-3 cm broad; stem 4-5 cm long, 2-3 mm thick.

Ground among fallen leaves in woods. Lewis co. September.

This is a rare species which departs from the generic character in its decurrent lamellae. It is unlike any of our other species of *Inocybe* in its minute globose spores. These have been described in *Sylloge* as echinulate, but in our plant the spores appear even under ordinary magnification, most minutely uneven under higher power or better definition. By some, the species has been referred to the genus *Flammula*. It has also been taken as the type species of a genus *Ripartites*, instituted by Karsten to include all the species of this section.

***Inocybe trechispora* (Berk.) Karst.**

ROUGH SPORE INOCYBE

Ag. (*Hebeloma*) *trechisporus* Berk. Outl. B. Fungi, p.156

Pileus thin, convex, acutely umbonate, at first viscid, then dry and silky, whitish with the umbo yellowish; lamellae subdistant, ventricose, sinuate, whitish becoming subferruginous; stem equal, slightly striate, stuffed, mealy, whitish; spores subglobose or ellipsoid, nodulose, 6-8 μ in diameter or 7-8 x 5-6 μ , cystidia 40-50 x 12-20 μ .

Pileus 2-3 cm broad; stem 2.5-5 cm long, 2-5 mm thick.

Ground in woods. Herkimer and Onondaga counties. August.

***Inocybe fuscodisca* (Pk.) Mass.**

BROWN DISK INOCYBE

Ag. (*Hebeloma*) *fuscodisca* Pk. N. Y. State Mus. Rep't 27, p.95, pl.1, fig.3-6

Pileus conic becoming campanulate or expanded, umbonate, slightly viscid, fibrillose, whitish, blackish brown on the umbo; lamellae close, adnexed, whitish becoming brownish ferruginous, white crenulate on the edge; stem equal, solid, whitish pruinose at the top, fibrillose below, brownish; spores ellipsoid, even, 8-10 x 5-6 μ , cystidia 40-70 x 12-20 μ .

Pileus 1.5-2.5 cm broad; stem 2.5-7 cm long, 2-4 mm thick.

Ground under trees. Sullivan co. September. Rare.

The viscid pellicle is separable. The odor resembles that of chestnut blossoms.

✓
Inocybe vatricosoides n. sp.

VATRICOSOID INOCYBE

Pileus thin, convex becoming nearly plane, slightly viscid when moist, obtuse or subumbonate, fibrillose on the margin from the abundant whitish webby veil, whitish, often reddish in the center, flesh whitish, odor like that of radishes; lamellae close, broadly sinuate, adnate with a decurrent tooth, whitish becoming brownish ferruginous, white crenulate on the edge; stem equal, flexuous usually curved at the base, stuffed or hollow, silky fibrillose, whitish or grayish, sometimes with whitish floccose scales toward the base; spores ellipsoid, even, $10-12 \times 6-8 \mu$.

Pileus 2-3 cm broad; stem 2.5-5 cm long, 2-6 mm thick.

Damp ground under willows. Ulster co. September.

This species is closely allied to *Inocybe vatricosa* Fr. to which it was referred in New York State Museum Report 41, page 67, but from which it is here separated because of its well developed webby veil, its radishlike odor, its adnate lamellae, its silky fibrillose stem and its larger spores.

Pileus tenuis, convexus, deinde subplanus, dum humidus subviscidus, obtusus subumbonatusve, margine velo abundante albido arachnoideo fibrillosus, albidus saepe centro rufescens, carne albida, odore Raphani; lamellae confertae, late sinuatae, adnatae, albidae deinde fusco-ferrugineae, acie albae crenulatae; stipes aequalis, flexuosus, saepe basi curvatus, farctus fistulosusve, sericeo-fibrillosus, aliquando infra squamulis albis floccosis ornatus, albidus vel griseus; sporae ellipsoideae, leves, $10-12 \times 6-8 \mu$.

Inocybe radiata Pk. N. Y. State Mus. Bul. 105, p. 24. The Port Jefferson specimens referred to this species as a small form are rather a small form of *Inocybe asterospora* Quel.

NEW YORK SPECIES OF HEBELOMA**Hebeloma** Fr.

Veil partial, fibrillose or obsolete; stem fleshy fibrous, somewhat mealy at the apex; margin of the pileus at first incurved, the cuticle continuous, glabrous, subviscid; lamellae sinuate, adnexed, usually whitish on the edge; spores subargillaceous. *Sylloge* 5:791

This genus formerly included the species now referred to *Inocybe*. It differs from it specially in its partial veil and in its continuous, subviscid and glabrous cuticle. Some of the species have a peculiar radishlike odor. The spores in all our species are even. As in the

preceding genus, most of the species are terrestrial. They have been placed in two primary divisions, Exannulata and Subannulata. Our species fall in the Exannulata division which has been divided into three sections, the principal characters of which are indicated in the following key.

KEY TO THE SECTIONS

- | | |
|------------------------------------|-----------|
| Pileus more than 3.5 cm broad..... | I |
| Pileus less than 3.5 cm broad..... | Pusilla |
| I Veil present..... | Indusiata |
| I Veil absent..... | Denudata |

Indusiata

Veil evident, webby, often making the margin of the pileus superficially silky.

KEY TO THE SPECIES

- | | |
|--|--------------|
| Pileus glutinous and squamose..... | glutinosum |
| Pileus viscid, not squamose..... | I |
| I Stem commonly showing fragments of an annulus..... | velatum |
| I Stem not at all annulate..... | 2 |
| 2 Pileus umbonate..... | firmum |
| 2 Pileus not umbonate..... | 3 |
| 3 Spores 6-8 long..... | parvifructum |
| 3 Spores 10-12 long..... | 4 |
| 4 Stem white, fibrous squamulose..... | fastibile |
| 4 Stem whitish or subochraceous, fibrillose..... | pascuense |

Hebeloma glutinosum (Lindgr.) Fr.

GLUTINOUS HEBELOMA

Sylloge 5:793

Pileus fleshy, convex becoming plane, glutinous, sprinkled with white superficial squamules, yellowish white, flesh white or whitish; lamellae close, sinuate, adnexed, yellowish becoming dingy cinnamon; stem equal or slightly thickened at the base, firm, stuffed, mealy at the top, fibrillose squamulose, whitish, somewhat ferruginous within; spores ellipsoid, 10-12 x 5-6 μ .

Pileus 2.5-7 cm broad; stem 4-8 cm long, 4-8 mm thick.

Ground in woods. Essex and Warren counties. September and October.

This species is easily recognized by its very viscose or glutinous pileus with its superficial white scales. These are not persistent and consequently specimens may be found without any scales.

Hebeloma velatum Pk.

VEILED HEBELOMA

Hebeloma colvini Pk., var. *velatum* Pk. N. Y. State Mus.
Rep't 48, p.19

Pileus convex, plane or slightly centrally depressed, obtuse or umbonate, slightly viscid when moist, glabrous or slightly silky from the veil which may disappear with age or persist and make the margin silky or floccosely scaly or appendiculate with its fragments, chestnut color, reddish gray, pale ochraceous or grayish; lamellae close, ventricose, adnexed, whitish becoming pale cinnamon, whitish and often crenulate on the edge; stem equal, hollow, silky fibrillose, sometimes floccosely squamulose toward the base, often more or less annulate, the soft cottony whitish or grayish veil rupturing and adhering partly to the stem and partly to the margin of the pileus, whitish; spores subellipsoid, $10-12 \times 6-8 \mu$.

Pileus 1.5-6 cm broad; stem 1.5-6 cm long, 4-6 mm thick.

Gregarious or cespitose. Gravelly soil under cottonwood trees. Clinton co. September.

This is an extremely variable species and shows how difficult it may be with a limited number of specimens in such cases to locate them correctly. All the forms here included under one name were collected at the same time and place, in a limited area but a few feet in diameter. They are without doubt all one species. Their general appearance suggested such a close relationship to *Hebeloma colvini* Pk. that it was thought best to group them all under that species as a variety distinguished chiefly by its more fully developed veil. If only the form having the veil and annulus in their most highly developed condition had been seen, the species might easily have been referred to the genus *Pholiota*. Even with those in which only fragmentary vestiges of the veil adhere to the stem its natural place would seem to be in the *Subannulata* division of the genus *Hebeloma*. But other forms show no trace of an annulus and compel us to be more conservative in our assignment of this perplexing species. It is therefore placed where the more abundant forms and less strongly developed or silky fibrillose veil would require it to go. It differs from *Hebeloma strophosum* Fr. in its great variability, differently colored pileus, radishlike odor and specially in the whitish color of the young lamellae.

Hebeloma firmum (Pers.) Fr.

FIRM HEBELOMA

Sylloge 5:793

Pileus fleshy, convex or campanulate becoming expanded, umbonate, viscid, fibrillose, brick red with paler margin; lamellae close, rounded behind, adnexed, tan color becoming subferruginous; stem equal or nearly so, solid or with a small cavity, floccosely squamulose, whitish sometimes becoming subferruginous toward the base; spores subellipsoid, $10-12 \times 5-6 \mu$.

Pileus 5-7 cm broad; stem 4-6 cm long, 6-8 mm thick.

Mossy ground in low woods: Essex co. September.

Hebeloma parvifructum Pk.

SMALL FRUIT HEBELOMA

Ag. (*Hebeloma*) *parvifructus* Pk. N. Y. State Mus. Rep't 38, p.88

Pileus convex becoming expanded, slightly viscid, whitish, grayish brown or pale chestnut, often paler on the margin; lamellae moderately close, slightly sinuate, white becoming dingy ochraceous, at first hidden by the copious white webby filaments of the veil; stem equal, often flexuous, solid, silky fibrillose, pruinose and substriate at the top, whitish above, ferruginous or brownish toward the base; spores subochraceous, $6-7 \times 4-5 \mu$.

Pileus 5-7 cm broad; stem 5-8 cm long, 6-8 mm thick.

Sandy soil in pine woods. Albany co. October.

The small spores are suggestive of the specific name.

Hebeloma fastibile Fr.

OCHERY HEBELOMA

Sylloge 5:792

Pileus convex or nearly plane, compact, often wavy, obtuse, viscid when young, whitish, yellowish or tan color, flesh white, odor similar to that of radishes, taste bitterish, veil webby, distinct; lamellae sinuate, adnexed, subdistant, whitish or pallid becoming cinnamon, whitish on the edge; stem equal, solid, fibrous, firm, sometimes slightly bulbous, white; spores ellipsoid, $10-12 \times 5-6 \mu$.

Pileus 3-7 cm broad; stem 5-7 cm long, 5-10 mm thick.

Ground in woods. Albany and Ulster counties. October.

A small white form, perhaps var. *alba* Sacc., has been found in Albany county.

Hebeloma pascuense Pk.

PASTURE HEBELOMA

N. Y. State Mus. Rep't 53, p.844, pl.C, fig.21-27

Pileus thin, convex becoming nearly plane, viscid when moist, obscurely innately fibrillose, brownish clay color, often darker or rufescent in the center, the margin when young often whitened by the thin webby veil, flesh whitish, odor similar to that of radishes; lamellae close, adnexed, whitish becoming pale ochraceous; stem firm, equal, solid, fibrillose, slightly mealy at the top, whitish or pallid; spores pale ochraceous, subellipsoid, uninucleate $10 \times 6 \mu$.

Pileus 2.5-5 cm broad; stem 2.5-5 cm long, 4-6 mm thick.

Gregarious or subcespitose. Stony pastures. Warren co. October.

Closely related to *Hebeloma fastibile* Fr. but a smaller species with a more slender stem, a different habit and habitat, differently colored pileus and more crowded lamellae. Sometimes a narrow brown zone or line encircles the pileus near the margin.

Denudata

Pileus glabrous, veil absent from the first.

The species are easily distinguished from those of the preceding section by the entire absence of a veil.

KEY TO THE SPECIES

- | | |
|---|-----------------|
| Pileus white or whitish..... | 1 |
| Pileus some other color..... | 3 |
| 1 Lamellae dingy flesh color..... | sarcophyllum |
| 1 Lamellae white or whitish becoming dingy ferruginous..... | 2 |
| 2 Pileus white or yellowish white..... | album |
| 2 Pileus whitish or grayish white..... | albidulum |
| 3 Plant having a radishlike odor..... | crustuliniforme |
| 3 Plant not having a radishlike odor..... | 4 |
| 4 Plant growing in sandy soil in open places..... | colvini |
| 4 Plant growing in woods..... | longicaudum |

Hebeloma sarcophyllum Pk.

PINK GILL HEBELOMA

Ag. (*Hebeloma*) *sarcophyllum* Pk. N. Y. State Cab. Rep't 23, p.96, pl.I, fig.7-11

Pileus fleshy, obtusely conic or convex, glabrous, white, flesh white, taste bitterish; lamellae subclose, adnexed, deeply sinuate, dingy flesh color; stem equal, firm, stuffed, mealy or minutely

squamulose at the top, white; spores dark ferruginous, subellipsoid, 8-10 x 5-6 μ .

Pileus 1.5-3 cm broad; stem 2.5-4 cm long, 2-4 mm thick.

Grassy ground. Rensselaer co. June. Rare.

A species well marked by the peculiar color of the lamellae which at first suggests a species of *Agaricus*.

***Hebeloma album* Pk.**

WHITE HEBELOMA

N. Y. State Mus. Rep't 54, p.147, pl.G, fig.1-7

Pileus fleshy, firm, convex becoming nearly plane or concave by the margin curving upward, glabrous, subviscid, white or yellowish white, flesh white; lamellae thin, narrow, close, sinuate, adnexed, whitish becoming brownish ferruginous; stem equal, firm, rather long, solid or stuffed, slightly mealy at the top, white; spores subellipsoid, pointed at one or both ends, 12-16 x 6-8 μ .

Pileus 2.5-5 cm broad; stem 3.5-7 cm long, 4-6 mm thick.

Among fallen leaves in woods. Essex co. October.

Easily recognized by the white color of both pileus and stem. A new figure of this species is given in the present report on plate 117, figure 1-6.

***Hebeloma albidulum* Pk.**

WHITISH HEBELOMA

N. Y. State Mus. Rep't 54, p.148

Pileus fleshy, firm, broadly convex or nearly plane, glabrous, slightly viscid when moist, dingy white or grayish white, flesh white; lamellae close, narrow, adnexed, whitish becoming brownish ferruginous, white and minutely denticulate on the edge; stem equal, firm, glabrous, slightly mealy or pruinose at the top, hollow, sometimes slightly bulbous, colored like the pileus; spores subellipsoid, obtuse, 10-12 x 6-8 μ .

Pileus 2.5-6 cm broad; stem 3-6 cm long, 4-6 mm thick.

Among fallen leaves in woods. Essex co. October.

This differs from *Hebeloma album* Pk. in its more dingy color, its hollow stem and its shorter and more obtuse spores.

Hebeloma crustuliniforme (Bull.) Fr.

CRUSTULINE HEBELOMA

Sylloge 5:799

Pileus fleshy, convex becoming plane, obtuse or with an obtuse umbo, even, glabrous, slightly viscid when young, whitish tan or brick red, odor like that of radishes; lamellae close, adnexed, narrow, thin, whitish becoming clay color or brownish ferruginous; stem equal, stuffed or hollow, subbulbous, white squamulose at the top, whitish; spores ellipsoid, unequal, $10-12 \times 5-7 \mu$.

Pileus 4-6 cm broad; stem 4-5 cm long, 6-10 mm thick.

Ground in woods or open places. Cattaraugus and Ulster counties. September.

Hebeloma colvini Pk.

COLVIN HEBELOMA

Ag. (*Hebeloma*) *colvini* Pk. N. Y. State Mus. Rep't 28, p.49

Pileus fleshy, convex or nearly plane, sometimes gibbous or broadly umbonate, rarely centrally depressed, glabrous, grayish or alutaceous with an ochraceous tint; lamellae close, broad, sinuate, adnexed, whitish becoming brownish ochraceous; stem equal, flexuous, silky fibrillose, stuffed or hollow above, solid toward the base, whitish; spores subellipsoid, $10-12 \times 5-6 \mu$.

Pileus 2.5-7.5 cm broad; stem 2.5-8 cm long, 2-6 mm thick.

Sandy soil in open places. Albany co. October.

The mycelium binds the sand into a globose mass which adheres to the base of the stem.

Hebeloma longicaudum (Pers.) Fr.

LONG STEM HEBELOMA

Sylloge 5:800

Pileus fleshy, convex becoming expanded, glabrous, viscid, whitish, argillaceous or tan color, sometimes brownish or yellowish red on the disk; lamellae close, sinuate near the stem, adnexed, whitish and serrulate on the edge, tan color becoming dingy cinnamon; stem unequal, rather long, fragile, partly hollow, mealy at the top, obsoletely fibrillose, white; spores oblong or ellipsoid, $10-12 \times 6-8 \mu$.

Pileus 3.5-6 cm broad; stem 5-9 cm long, 5-8 mm thick.

Ground in woods. Catskill mountains. September.

A rare species in our State.

Pusilla

Pileus small, less than 3.5 cm broad.

Several species have been referred to this section that have not a viscid pileus, but in some it is moist.

KEY TO THE SPECIES

- Pileus slightly viscid when moist.....1
- Pileus not viscid when moist.....4
- 1 Pileus white or yellowish white.....sociale
- 1 Pileus some other color.....2
- 2 Pileus ochraceous yellow.....gregarium
- 2 Pileus tawny brown or reddish brown.....3
- 3 Stem 2.5 cm long, center of pileus not changing color.....sordidulum
- 3 Stem longer, center of pileus changing color with age
or in drying.....discomorbidum
- 4 Pileus hygrophanous when moist.....5
- 4 Pileus not hygrophanous when moist.....7
- 5 Stem white.....palustre
- 5 Stem not white.....6
- 6 Pileus uniformly brown.....illicitum
- 6 Pileus brown with a whitish or pallid margin.....pallidomarginatum
- 7 Pileus broadly umbonate, stem solid.....excedens
- 7 Pileus not umbonate, stem hollow.....fragilius

Hebeloma sociale Pk.

SOCIAL HEBELOMA

N. Y. State Mus. Bul. 75, p.15

Pileus fleshy but thin, convex becoming plane or nearly so, glabrous, slightly viscid when moist, yellowish white, flesh yellowish white, taste nauseous; lamellae thin, close, adnexed, whitish, then yellowish, finally brownish ferruginous; stem short, fibrous, floc-cose fibrillose, hollow, white; spores brownish ferruginous, ellip-soid, 6-8 x 4-5 μ .

Pileus 2-3 cm broad; stem 2.5-3.5 cm long, 3-6 mm thick.

Gregarious or subcespitose. Among short grass in pastures. Albany co. October.

This is distinguished from our other white or whitish species by its peculiar habitat and its small spores.

Hebeloma gregarium Pk.

GREGARIOUS HEBELOMA

N. Y. State Mus. Rep't 49, p.18

Pileus thin, hemispheric or convex, obtuse or rarely with a small inconspicuous umbo, slightly viscid when moist, glabrous or slightly silky on the margin, pale ochraceous, sometimes with a reddish or tawny tint in the center, flesh whitish; lamellae thin, close, adnate, whitish becoming subcinnamon; stem slender, stuffed or hollow, fibrillose, whitish, slightly mealy or pruinose at the top; spores ellipsoid, $10-11 \times 5-6 \mu$.

Pileus 2-3.5 cm broad; stem 3-5 cm long, 2-4 mm thick.

Sandy soil in heathy places. Albany co. October. Rare.

The pileus is sometimes split on the margin in such a way as to cause it to appear stellately lobed. A slight radishlike odor is perceptible when the pileus is cut or broken. The plants are gregarious.

Hebeloma sordidulum Pk.

SLIGHTLY SORDID HEBELOMA

Ag. (*Hebeloma*) *sordidulus* Pk. N. Y. State Mus. Rep't 38, p.88

Pileus thin, firm, viscid when moist, convex, brownish red or tawny brown, paler on the margin, flesh white, with a radishlike odor; lamellae broad, close, rounded behind, slightly adnexed, pallid becoming brownish ochraceous; stem short, equal, stuffed or hollow, slightly fibrillose, pruinose at the top, white; spores sub-ellipsoid, $12-15 \times 6-7 \mu$.

Pileus 2-4 cm broad; stem about 2.5 cm long, 3-4 mm thick.

Sandy soil in open bushy places. Albany co. October. Rare.

Hebeloma discomorbidum Pk.

DISK DISEASED HEBELOMA

Ag. (*Naucoria*) *discomorbidus* Pk. N. Y. State Mus. Rep't 26, p.58

Pileus thin, broadly convex or nearly plane, glabrous, slightly viscid, reddish brown or chestnut color becoming brown in the center with age or in drying; lamellae close, narrow, white or pallid becoming brownish ferruginous, white and crenulate on the edge; stem equal, stuffed or hollow, slightly mealy at the top, white; spores ellipsoid, uninucleate, $10 \times 6 \mu$.

Pileus 2-4 cm broad; stem 4-6 cm long, 2-4 mm thick.

Ground in woods. Lewis and Columbia counties. September and October.

In the dried specimens the center of the pileus has a brown or discolored appearance as if beginning to decay. This is suggestive of the specific name.

Hebeloma palustre Pk.

MARSH HEBELOMA

N. Y. State Mus. Bul. 25, p.649

Pileus thin, broadly convex becoming nearly plane, sometimes wavy or irregular, glabrous, hygrophanous, grayish brown and slightly striatulate on the margin when moist, paler when dry, flesh whitish; lamellae close, thin, ventricose, adnexed, grayish white becoming brownish cinnamon; stem equal or tapering upward, hollow, silky, white; spores subellipsoid, uninucleate, $10-12 \times 6-8 \mu$.

Pileus 2.5-3.5 cm broad; stem 5-7 cm long, 4-8 mm thick.

Mossy ground in swampy woods. Oswego co. October. Rare.

Hebeloma illicitum Pk.

UNLAWFUL HEBELOMA

Ag. (*Hebeloma*) *illicitus* Pk. N. Y. State Mus. Rep't 24, p.68, pl.4, fig.1-5

Pileus fleshy, firm, convex or expanded, obtuse, glabrous, hygrophanous, dark brown when moist, paler when dry; lamellae close, broad, ventricose, adnexed, pale brown; stem equal, firm, hollow, striate at the top, with a white mycelium at the base, colored like but paler than the pileus; spores subellipsoid, $8-10 \times 4-5 \mu$.

Pileus 2.5-3.5 cm broad; stem 3.5-5 cm long, 4 mm thick.

Decaying wood and sticks in woods. Lewis and Washington counties. September. Rare.

Gregarious or caespitose. The specific name has reference to its habitat, which is unusual for species of this genus.

Hebeloma pallidomarginatum Pk.

PALE MARGIN HEBELOMA

Ag. (*Hebeloma*) *pallidomarginatus* Pk. N. Y. State Mus. Rep't 25, p.78

Pileus broadly convex, sometimes irregular, glabrous, hygrophanous, brown with a pale margin when moist, ochraceous and subatomaceous when dry; lamellae close, thin, adnexed, brownish ochraceous; stem commonly long and flexuous, equal or tapering upward, hollow, white floccose at the base, colored like but paler than the pileus; spores subellipsoid, $10 \times 5 \mu$.

Pileus 1-2.5 cm broad; stem 2.5-7 cm long, 2 mm thick.

Gregarious in swamps and wet places. Rensselaer co. September. Rare.

Hebeloma excedens Pk.

THIN MARGIN HEBELOMA

Ag. (*Hebeloma*) *excedens* Pk. N. Y. State Mus. Rep't 24, p.68

Pileus thin, convex, obtuse or broadly umbonate, glabrous, pale alutaceous, the margin surpassing the lamellae, taste and odor like that of radishes; lamellae close, deeply sinuate, adnexed, moderately broad, minutely eroded on the edge, pallid becoming brownish ferruginous; stem equal, solid, silky fibrillose, colored like the pileus; spores subellipsoid, $10-12 \times 6-7 \mu$.

Pileus 1.5-2.5 cm broad; stem 3-5 cm long, 2-4 mm thick.

Sandy soil under or near pine trees. Saratoga co. October. Very rare. Not found since 1870.

Easily known by its thin margin which extends beyond the lamellae.

Hebeloma fragilius Pk.

FRAGILE HEBELOMA

Ag. (*Hebeloma*) *fragilior* Pk. N. Y. State Mus. Rep't 27, p.95

Pileus thin, fragile, convex becoming plane or centrally depressed, sometimes irregular or wavy on the margin, minutely squamulose when young, soon glabrous, pale grayish ochraceous; lamellae subdistant, ventricose, adnexed, whitish and crenulate on the edge, subochraceous; stem slender, equal, hollow, minutely furfuraceous becoming glabrous, colored like the pileus; spores ellipsoid, $6 \times 4 \mu$.

Pileus 6-12 mm broad; stem about 2.5 cm long, 1-2 mm thick.

Damp decaying leaves in wet places in swamps. Hamilton co. July. Rare.

This is a very small species and in habitat is unlike any other of our species. Sometimes the stem is expanded at the base in a thin disk closely applied to the matrix. It needs further investigation and may possibly be found to be a species of *Naucoria*.

Ag. (*Hebeloma*) *lacerus* Fr. N. Y. State Cab. Rep't 23, p. 95 is referable to *Inocybe asterospora* Quel.

Ag. (*Hebeloma*) *flocculosus* Berk. N. Y. State Cab. Rept 23, p. 96 is referable to *Inocybe infelix* Pk.

Ag. (*Hebeloma*) *ascophorus* Pk. N. Y. State Mus. Rep't 24, p. 68 is erroneously described and is referable to *Flammula highlandensis* Pk.

LIST OF EDIBLE, POISONOUS AND UNWHOLESOME
MUSHROOMS HITHERTO FIGURED AND DESCRIBED
BY C. H. PECK, STATE BOTANIST

Agaricus abruptus *Pk.*

N. Y. State Mus. Mem. 4, p.163-64, pl.59, fig.8-14. 1900. (*Agaricus abruptibulbus* *Pk.* N. Y. State Mus. Bul. 94, p.36. 1905)

Agaricus arvensis *Schaeff.*

N. Y. State Mus. Rep't 48, p.140-41, pl.8. 1896. Bot. ed.

Agaricus campester *L.*

N. Y. State Mus. Rep't 48, p.134-37, pl.6. 1896. Bot. ed.

Agaricus diminutivus *Pk.*

N. Y. State Mus. Rep't 54, p.184-85, pl.74, fig.1-8. 1901

Agaricus haemorrhoidarius *Schulz.*

N. Y. State Mus. Rep't 54, p.183-84, pl.75. 1901

Agaricus micromegethus *Pk.*

(*Agaricus pusillus* *Pk.* N. Y. State Mus. Rep't 54, p.152. 1901)

N. Y. State Mus. Bul. 116, p.44, pl.107, fig.1-6. 1907

Agaricus placomyces *Pk.*

N. Y. State Mus. Rep't 48, p.142-43, pl.9, fig.7-12. 1896. Bot. ed.

Agaricus rodmani *Pk.*

N. Y. State Mus. Rep't 48, p.137-38, pl.9, fig.1-6. 1896. Bot. ed.

Agaricus silvicola *Pk.*

N. Y. State Mus. Mem. 4, p.164-65, pl.59, fig.1-7. 1900

Agaricus subrufescens *Pk.*

N. Y. State Mus. Rep't 48, p.138-40, pl.7. 1896. Bot. ed.

Amanita caesarea *Scop.*

N. Y. State Mus. Rep't 48, p.155-57, pl.15. 1896. Bot. ed.

Amanita rubescens *Fr.*

N. Y. State Mus. Rep't 48, p.157-59, pl.16. 1896. Bot. ed.

Amanitopsis strangulata (*Fr.*) *Roze*

N. Y. State Mus. Rep't 51, p.300-2, pl.50, fig.1-10. 1898

N. Y. State Mus. Mem. 4, p.134-35, pl.44, fig.1-10. 1900

Amanitopsis vaginata *Roze*

N. Y. State Mus. Rep't 48, p.159-60, pl.17. 1896. Bot. ed.

Armillaria mellea *Vahl*

N. Y. State Mus. Rep't 48, p.164-67, pl.20. 1896. Bot. ed.

Boletinus grisellus *Pk.*

N. Y. State Mus. Mem. 4, p.169, pl.52, fig.13-19. 1900

Boletinus pictus *Pk.*

N. Y. State Mus. Bul. 25, p.681-82, pl.61, fig.1-5. 1899

N. Y. State Mus. Mem. 4, p.169, pl.61, fig.1-5. 1900

Boletus affinis *Pk.*

N. Y. State Mus. Rep't 49, p.64, pl.48, fig.6-16. 1896. Bot. ed.

N. Y. State Mus. Mem. 4, p.174-75, pl.66, fig.7-14. 1900

Boletus bicolor *Pk.*

N. Y. State Mus. Bul. 54, p.973-74, pl.81, fig.6-11. 1902

Boletus brevipes *Pk.*

N. Y. State Mus. Rep't 49, p.63-64, pl. 48, fig.1-5. 1896. Bot. ed.

N. Y. State Mus. Mem. 4, p.174, pl.66, fig.1-6. 1900

Boletus castaneus *Bull.*

N. Y. State Mus. Rep't 48, p.201-2, pl.36, fig.1-7. 1895. Bot. ed.

Boletus chrysenteron albocarneus *Pk.*

N. Y. State Mus. Rep't 54, p.185-86, pl.76, fig.21-25. 1901

Boletus clintonianus *Pk.*

N. Y. State Mus. Bul. 25, p.682, pl.61, fig.6-10. 1899

N. Y. State Mus. Mem. 4, p.170-71, pl.63. 1900

Boletus edulis *Bull.*

N. Y. State Mus. Rep't 47, p.200-1, pl.36, fig.8-12. 1896. Bot. ed.

Boletus edulis clavipes *Pk.*

N. Y. State Mus. Rep't 51, p.309-10, pl.54. 1898

N. Y. State Mus. Mem. 4, p.173-74, pl.65. 1900

Boletus eximius *Pk.*

N. Y. State Mus. Bul. 54, p.976-77, pl.80, fig.6-12. 1902

Boletus frostii *Russell*

N. Y. State Mus. Bul. 116, p.44-45, pl.108. 1907

Boletus granulatus *L.*

N. Y. State Mus. Rep't 48, p.196-97, pl.34, fig.1-5. 1896. Bot. ed.

Boletus laricinus *Berk.*

N. Y. State Mus. Bul. 94, p.46-47, pl.89. 1905

Boletus luteus *L.*

N. Y. State Mus. Rep't 48, p.195-96, pl.33, fig.7-12. 1896. Bot. ed.

Boletus niveus *Fr.*

N. Y. State Mus. Bul. 122, p.140-41, pl.113. 1908

Boletus nobilis *Pk.*

N. Y. State Mus. Bul. 94, p.48, pl.91. 1905

Boletus ornatipes *Pk.*

N. Y. State Mus. Bul. 54, p.975-76, pl.80, fig.1-5. 1902

Boletus pallidus *Frost*

N. Y. State Mus. Bul. 54, p.974-75, pl.81, fig.1-5. 1902

Boletus rubropunctus *Pk.*

N. Y. State Mus. Bul. 94, p.47, pl.90. 1905

Boletus rugosiceps *Pk.*

N. Y. State Mus. Bul. 116, p.45. 1907

N. Y. State Mus. Bul. 94, p.20-21, pl.Q, fig.6-10. 1905

Boletus scaber *Fr.*

N. Y. State Mus. Rep't 48, p.199-200, pl.35. 1896. Bot. ed.

Boletus spectabilis *Pk.*

N. Y. State Mus. Mem. 4, p.171-72, pl.62. 1900

Boletus subaureus *Pk.*

N. Y. State Mus. Mem. 4, p.169-70, pl.61, fig.6-13. 1900

Boletus subglabripes *Pk.*

N. Y. State Mus. Rep't 51, p.308-9, pl.55. 1898

N. Y. State Mus. Mem. 4, p.172-73, pl.64. 1900

Boletus subluteus *Pk.*

N. Y. State Mus. Rep't 48, p.196, pl.33, fig.1-6. 1896. Bot. ed.

Boletus versipellis Fr.

N. Y. State Mus. Rep't 48, p.198, pl.34, fig.6-10. 1896. Bot. ed.

Bovista pila B. & C.

N. Y. State Mus. Bul. 75, p.34, pl.84, fig.14-18. 1904

Bovista plumbea Pers.

N. Y. State Mus. Bul. 54, p.977-78, pl.81, fig.12-19. 1902

Cantharellus cibarius Fr.

N. Y. State Mus. Rep't 48, p.190-91, pl.32. 1896. Bot. ed.

Cantharellus cinnabarinus Schw.

N. Y. State Mus. Bul. 25, p. 679-680, pl.60, fig.1-9. 1899

N. Y. State Mus. Mem. 4, p.155-56, pl.55, fig.1-8. 1900

Cantharellus dichotomus Pk.

N. Y. State Mus. Bul. 67, p.46-47, pl.84, fig.8-21. 1903

Cantharellus floccosus Schw.

N. Y. State Mus. Bul. 25, p.680-81, pl.60, fig.10-14. 1899

N. Y. State Mus. Mem. 4, p.156-57, pl.55, fig.9-13. 1900

Cantharellus infundibuliformis (Scop.) Fr.

N. Y. State Mus. Mem. 4, p.158-59, pl.56, fig.9-16. 1900

Cantharellus lutescens Fr.

N. Y. State Mus. Mem. 4, p.157-58, pl.56, fig.1-8. 1900

Cantharellus minor Pk.

N. Y. State Mus. Bul. 131, p.41-42, pl.116, fig.12-17. 1909

Clavaria botrytis Pers.

N. Y. State Mus. Rep't 48, p.211, pl.39, fig.5-7. 1896. Bot. ed.

Clavaria botrytoides Pk.

N. Y. State Mus. Bul. 94, p.49, pl.93, fig.5-7. 1905

Clavaria conjuncta Pk.

N. Y. State Mus. Bul. 105, p.42-43, pl.102. 1906

Clavaria cristata Pers.

N. Y. State Mus. Rep't 48, p.211-12, pl.39, fig.8-12. 1896. Bot. ed.

Clavaria flava Schaeff.

N. Y. State Mus. Rep't 48, p.210, pl.39, fig.1-4. 1896. Bot. ed.

Clavaria pistillaris L.

N. Y. State Mus. Bul. 94, p.50, pl.93, fig.1-4. 1905

Clavaria pistillaris umbonata Pk.

N. Y. State Mus. Mem. 4, p.178, pl.66, fig.15-17. 1900

Clitocybe adirondackensis Pk.

N. Y. State Mus. Rep't 54, p.174-75, pl.69, fig.1-13. 1901

Clitocybe amethystina (Bolt.) Pk.

N. Y. State Mus. Bul. 116, p.40-41, pl.106, fig.1-6. 1907

Clitocybe clavipes (Pers.) Fr.

N. Y. State Mus. Rep't 49, p.58, pl.45, fig.1-7. 1896. Bot. ed.

N. Y. State Mus. Mem. 4, p.139-40, pl.46, fig.1-6. 1900

Clitocybe infundibuliformis Schaeff.

N. Y. State Mus. Rep't 48, p.174-75, pl.24, fig.1-6. 1896. Bot. ed.

Clitocybe laccata Scop.

N. Y. State Mus. Rep't 48, p.175-77, pl.25. 1896. Bot. ed.

Clitocybe maculosa Pk.

N. Y. State Mus. Rep't 54, p.174, pl.69, fig.14-21. 1901

Clitocybe media *Pk.*

N. Y. State Mus. Rep't 48, p.173-74, pl.23, fig.1-7. 1896. Bot. ed.

Clitocybe monadelpha *Morg.*

N. Y. State Mus. Rep't 51, p.302-3, pl.51, fig.1-5. 1898

N. Y. State Mus. Mem. 4, p.140-41, pl.46, fig.7-12. 1900

Clitocybe multififormis *Pk.*

N. Y. State Mus. Mem. 4, p.141, pl.47, fig.1-9. 1900

Clitocybe nebularis *Batsch*

N. Y. State Mus. Rep't 48, p.172-73, pl.23, fig.8-13. 1896. Bot. ed.

Clitocybe ochropurpurea *Berk.*

N. Y. State Mus. Bul. 116, p.41-42, pl.106, fig.7-11. 1907

Clitocybe subcylathiformis *Pk.*

N. Y. State Mus. Bul. 122, p.136-37, pl.110, fig.1-6. 1908

Clitopilus abortivus *B. & C.*

N. Y. State Mus. Bul. 54, p.968-69, pl.78, fig.13-19. 1902

Clitopilus micropus *Pk.*

N. Y. State Mus. Bul. 54, p.970, pl.78, fig.1-12. 1902

Clitopilus orcella *Bull.*

N. Y. State Mus. Rep't 48, p.153, pl.14, fig.7-11. 1896. Bot. ed.

Clitopilus prunulus *Scop.*

N. Y. State Mus. Rep't 48, p.152-53, pl.14, fig.1-6. 1896. Bot. ed.

Collybia acervata *Fr.*

N. Y. State Mus. Bul. 75, p.27-28, pl.84, fig.8-13. 1904

Collybia dryophila (*Bull.*) *Fr.*

N. Y. State Mus. Bul. 122, p.137-38, pl.111. 1908

Collybia familia *Pk.*

N. Y. State Mus. Bul. 75, p.28-29, pl.84, fig.1-7. 1904

Collybia platyphylla *Fr.*

N. Y. State Mus. Mem. 4, p.142-43, pl. 49. 1900

Collybia radicata (*Relh.*) *Fr.*

N. Y. State Mus. Rep't 51, p.304-5, pl.52. 1898

N. Y. State Mus. Mem. 4, p.143-44, pl.48. 1900

Collybia velutipes (*Curt.*) *Fr.*

N. Y. State Mus. Rep't 51, p.305-6, pl.50, fig.11-16. 1898

N. Y. State Mus. Mem. 4, p.144-45, pl.47, fig.10-15. 1900

Coprinus atramentarius *Fr.*

N. Y. State Mus. Rep't 48, p.144-45, pl.11, fig.7-11. 1896. Bot. ed.

Coprinus comatus *Fr.*

N. Y. State Mus. Rep't 48, p.143-44, pl.10. 1896. Bot. ed.

Coprinus micaceus *Fr.*

N. Y. State Mus. Rep't 48, p.145-47, pl.11, fig.1-6. 1896. Bot. ed.

Cortinarius cinnamomeus *Fr.*

N. Y. State Mus. Rep't 48, p.149-50, pl.13, fig.7-20. 1896. Bot. ed.

Cortinarius collinitus *Fr.*

N. Y. State Mus. Rep't 48, p.149, pl.13, fig.1-6. 1896. Bot. ed.

Cortinarius corrugatus *Pk.*

N. Y. State Mus. Bul. 25, p.674, pl.57, fig.6-13. 1899

N. Y. State Mus. Mem. 4, p.161-62, pl.58, fig.8-15. 1900

Cortinarius evernius *Fr.*

N. Y. State Mus. Mem. 4, p.162-63, pl.58, fig.1-7. 1900

Cortinarius violaceus Fr.

N. Y. State Mus. Rep't 48, p.148-49, pl.12. 1896. Bot. ed.

Craterellus cantharellus (Schw.) Fr.

N. Y. State Mus. Rep't 49, pl. 44, fig. 1-5. 1896. Bot. ed.

N. Y. State Mus. Mem. 4, p.177-78, pl.56, fig.17-21. 1900

Craterellus cornucopioides Pers.

N. Y. State Mus. Rep't 48, p.208-9, pl.24, fig.7-10. 1896. Bot. ed.

Crepidotus malachius B. & C.

N. Y. State Mus. Bul. 122, p.139, pl.112, fig.1-4. 1908

Fistulina hepatica Fr.

N. Y. State Mus. Rep't 48, p.204-5, pl.37, fig.5-9. 1896. Bot. ed.

Gyromitra esculenta Fr.

N. Y. State Mus. Rep't 48, p.128-29, pl.5, fig.1-3. 1896. Bot. ed.

Helvella crispa Fr.

N. Y. State Mus. Rep't 48, p.129-30, pl.5, fig.4-7. 1896. Bot. ed.

Hydnum albidum Pk.

N. Y. State Mus. Rep't 51, p.310, pl.56, fig.1-7. 1898

N. Y. State Mus. Mem. 4, p.175-76, pl.67, fig.1-7. 1900

Hydnum caput-ursi Fr.

N. Y. State Mus. Rep't 51, p.310-12, pl.56, fig.8-12. 1898

N. Y. State Mus. Mem. 4, p.176-77, pl.67, fig.8-12. 1900

Hydnum coralloides Scop.

N. Y. State Mus. Rep't 48, p.207-8, pl.24, fig.7-10. 1896. Bot. ed.

Hydnum repandum L.

N. Y. State Mus. Rep't 48, p.206-7, pl.38. 1896. Bot. ed.

Hygrophorus cantharellus Schw.

N. Y. State Mus. Rep't 54, p.175-76, pl.76, fig.8-20. 1901

Hygrophorus chlorophanus Fr.

N. Y. State Mus. Mem. 4, p.147, pl.51, fig.13-20. 1900

Hygrophorus flavodiscus Frost

N. Y. State Mus. Rep't 51, p.303-4, pl.51, fig.6-11. 1898

N. Y. State Mus. Mem. 4, p.145, pl.50, fig.1-6. 1900

Hygrophorus fuliginosus Frost

N. Y. State Mus. Rep't 49, p.59, pl.45, fig.8-14. 1896. Bot. ed.

N. Y. State Mus. Mem. 4, p.146, pl.50, fig.7-12. 1900

Hygrophorus laricinus Pk.

N. Y. State Mus. Mem. 4, p.146-47, pl.51, fig.1-12. 1900

Hygrophorus lauræ Morg.

N. Y. State Mus. Bul. 54, p.967-68, pl.77, fig.6-14. 1902

Hygrophorus lauræ decipiens Pk.

N. Y. State Mus. Bul. 94, p.46, pl.88, fig.8-11. 1905

Hygrophorus miniatus Fr.

N. Y. State Mus. Rep't 48, p.182-84, pl.28, fig.1-10. 1896. Bot. ed.

Hygrophorus nitidus B. & C.

N. Y. State Mus. Bul. 94, p.45, pl.88, fig.1-7. 1905

Hygrophorus pratensis Fr.

N. Y. State Mus. Rep't 48, p.181-82, pl.28, fig.11-17. 1896. Bot. ed.

Hygrophorus pudorinus Fr.

N. Y. State Mus. Bul. 67, p.41-42, pl.83, fig.1-6. 1903

Hygrophorus puniceus Fr.

N. Y. State Mus. Bul. 25, p.675, pl.58, fig.1-7. 1899

N. Y. State Mus. Mem. 4, p.149, pl.52, fig.1-7. 1900

Hygrophorus speciosus Pk.

N. Y. State Mus. Mem. 4, p.148, pl.51, fig.21-28. 1900

Hygrophorus virgineus (Wulf.) Fr.

N. Y. State Mus. Bul. 25, p.675-76, pl.58, fig.8-12. 1899

N. Y. State Mus. Mem. 4, p.150, pl.52, fig.8-12. 1900

Hypholoma aggregatum sericeum Pk.

N. Y. State Mus. Bul. 54, p.972-73, pl.79, fig.8-14. 1902

Hypholoma incertum Pk.

N. Y. State Mus. Bul. 25, p.676-77, pl.58, fig.13-20. 1899

N. Y. State Mus. Mem. 4, p.165-66, pl.60, fig.1-9. 1900

Hypholoma perplexum Pk.

N. Y. State Mus. Rep't 49, p.61-62, pl.47, fig.11-18. 1896. Bot. ed.

N. Y. State Mus. Mem. 4, p.166-67, pl.60, fig.10-17. 1900

Hypomyces lactifluorum (Schw.) Tul.

N. Y. State Mus. Bul. 105, p.43-44, pl.103. 1906

Lactarius chelidonium Pk.

N. Y. State Mus. Bul. 25, p.677-78, pl.59, fig.1-6. 1899

N. Y. State Mus. Mem. 4, p. 150-51, pl.53, fig.1-6. 1900

Lactarius deceptivus Pk.

N. Y. State Mus. Rep't 54, p.177-78, pl.70, fig.7-11. 1901

Lactarius deliciosus Fr.

N. Y. State Mus. Rep't 48, p.185-86, pl.29. 1896. Bot. ed.

Lactarius distans Pk.

N. Y. State Mus. Bul. 25, p.678-79, pl.59, fig.7-11. 1899

N. Y. State Mus. Mem. 4, p.151-52, pl.53, fig.7-11. 1900

Lactarius gerardii Pk.

N. Y. State Mus. Bul. 25, p.679, pl.59, fig.12-16. 1899

N. Y. State Mus. Mem. 4, p.152-53, pl.53, fig.12-16. 1900

Lactarius luteolus Pk.

N. Y. State Mus. Bul. 67, p.43, pl.83, fig.7-11. 1903

Lactarius rimosellus Pk.

N. Y. State Mus. Bul. 105, p.37, pl.95, fig.1-6. 1906

Lactarius serifluus (DC.) Fr.

N. Y. State Mus. Bul. 105, p.37-38, pl.95, fig.7-11. 1906

Lactarius subdulcis (Bull.) Fr.

N. Y. State Mus. Bul. 67, p.43-45, pl.83, fig.12-24. 1903

Lactarius subpurpureus Pk.

N. Y. State Mus. Rep't 54, p.176-77, pl.70, fig.1-6. 1901

Lactarius volemus Fr.

N. Y. State Mus. Rep't 48, p.186-88, pl.30. 1896. Bot. ed.

Lepiota americana Pk.

N. Y. State Mus. Rep't 49, p.56-57, pl.44, fig.6-10. 1896. Bot. ed.

N. Y. State Mus. Mem. 4, p.136-37, pl.44, fig.11-16. 1900

Lepiota cepaestipes Sow.

N. Y. State Mus. Bul. 94, p.44-45, pl.87. 1905

Lepiota clypeolaria (Bull.) Fr.

N. Y. State Mus. Rep't 54, p.173, pl.76, fig.1-7. 1901

Lepiota naucinoides *Pk.*

N. Y. State Mus. Rep't 48, p.162-64, pl.19. 1896. Bot. ed.

Lepiota procera *Scop.*

N. Y. State Mus. Rep't 48, p.161-62, pl.18. 1896. Bot. ed.

Lycoperdon cyathiforme *Bosc*

N. Y. State Mus. Rep't 48, p.121-22, pl.2. 1896. Bot. ed.

Lycoperdon gemmatum *Batsch*

N. Y. State Mus. Bul. 122, p.135-36, pl.114, fig.7-15. 1908

Lycoperdon giganteum *Batsch*

N. Y. State Mus. Rep't 48, p.121, pl.1. 1896. Bot. ed.

Lycoperdon subincarnatum *Pk.*

N. Y. State Mus. Bul. 122, p.135, pl.114, fig.1-6. 1908

Marasmius oreades *Fr.*

N. Y. State Mus. Rep't 48, p.195-96, pl.33, fig.7-12. 1896. Bot. ed.

Mitrula vitellina irregularis *Pk.*

N. Y. State Mus. Rep't 48, p.130-31, pl.5, fig.8-14. 1896. Bot. ed.

Morchella angusticeps *Pk.*

N. Y. State Mus. Rep't 48, p.125, pl.4, fig.5-9. 1896. Bot. ed.

Morchella bispora *Sor.*

N. Y. State Mus. Rep't 48, p.126-27, pl.3, fig.8-10. 1896. Bot. ed.

Morchella conica *Pers.*

N. Y. State Mus. Rep't 48, p.124-25, pl.4, fig.1-4. 1896. Bot. ed.

Morchella deliciosa *Fr.*

N. Y. State Mus. Rep't 48, p.125-26, pl.3, fig.4-7. 1896. Bot. ed.

Morchella esculenta *Pers.*

N. Y. State Mus. Rep't 48, p.124, pl.3, fig.1-3. 1896. Bot. ed.

Morchella semilibera *DC.*

N. Y. State Mus. Rep't 48, p.126, pl.3, fig.11-13. 1896. Bot. ed.

Paxillus involutus *Fr.*

N. Y. State Mus. Rep't 48, p.150-51, pl.28, fig.18-23. 1896. Bot. ed.

Pholiota adiposa *Fr.*

N. Y. State Mus. Rep't 49, p.60-61, pl.46, fig.18-23. 1896. Bot. ed.

N. Y. State Mus. Mem. 4, p.160-61, pl.57, fig. 12-17. 1900

Pholiota caperata *Pers.*

N. Y. State Mus. Rep't 54, p.182, pl.73, fig.1-5. 1901

Pholiota duroides *Pk.*

N. Y. State Mus. Bul. 131, p.39-40, pl.116, fig.1-7. 1909

Pholiota praecox (*Pers.*) *Fr.*

N. Y. State Mus. Rep't 49, p.59-60, pl.46, fig.1-17. 1896. Bot. ed.

N. Y. State Mus. Mem. 4, p.159-60, pl.57, fig.1-11. 1900

Pholiota squarrosa *Muell.*

N. Y. State Mus. Bul. 54, p.971-72, pl.79, fig.1-7. 1902

Pholiota squarrosoides *Pk.*

N. Y. State Mus. Rep't 54, p.183, pl.73, fig.6-15. 1901

Pholiota vermiflua *Pk.*

N. Y. State Mus. Bul. 75, p.32, pl.86, fig.12-20. 1904

Phylloporus rhodoxanthus (*Schw.*) *Bres.*

N. Y. State Mus. Bul. 131, p.40-41, pl.116, fig.8-11. 1909

Pleurotus ostreatus *Fr.*

N. Y. State Mus. Rep't 48, p.180-81, pl.26, fig.5-9. 1896. Bot. ed.

Pleurotus sapidus *Kalchb.*

N. Y. State Mus. Rep't 48, p.179-80, pl.27. 1896. Bot. ed.

Pleurotus ulmarius *Bull.*

N. Y. State Mus. Rep't 48, p.177-79, pl.26, fig.1-4. 1896. Bot. ed.

Pluteus cervinus (*Schaeff.*) *Fr.*

N. Y. State Mus. Rep't 54, p.181-82, pl.74, fig.9-19. 1901

Polyporus sulphureus *Fr.*

N. Y. State Mus. Rep't 48, p.203-4, pl.37, fig.1-4. 1896. Bot. ed.

Psilocybe foenicicii (*Pers.*) *Fr.*

N. Y. State Mus. Bul. 75, p.33-34, pl.86, fig.1-11. 1904

Russula abietina *Pk.*

N. Y. State Mus. Rep't 54, p.180-81, pl.72, fig.1-11. 1901

Russula albida *Pk.*

N. Y. State Mus. Bul. 105, p.38, pl.96. 1906

Russula brevipes *Pk.*

N. Y. State Mus. Rep't 54, p.178-79, pl.71, fig.1-5. 1901

Russula compacta *Frost*

N. Y. State Mus. Bul. 116, p.42, pl.109. 1907

Russula crustosa *Pk.*

N. Y. State Mus. Bul. 67, p.45-46, pl.84, fig.1-7. 1903

Russula earlei *Pk.*

N. Y. State Mus. Bul. 116, p.42. 1907

N. Y. State Mus. Bul. 67, p.24, pl.N, fig.5-10. 1903

Russula flavida *Frost*

N. Y. State Mus. Bul. 105, p.38-39, pl.97. 1906

Russula furcata (*Pers.*) *Fr.*

N. Y. State Mus. Bul. 75, p.31-32, pl.85, fig.9-14. 1904

Russula mariae *Pk.*

N. Y. State Mus. Bul. 75, p.29-31, pl.85, fig.1-8. 1904

Russula nigricans (*Bull.*) *Fr.*

N. Y. State Mus. Rep't 54, p.178, pl.71, fig.6-9. 1901

Russula ochrophylla *Pk.*

N. Y. State Mus. Rep't 51, p.307-8, pl.53, fig.8-14. 1898

N. Y. State Mus. Mem. 4, p.154-55, pl.54, fig.8-14. 1900

Russula pectinatoides *Pk.*

N. Y. State Mus. Bul. 116, p.43, pl.105, fig.6-10. 1907

Russula pusilla *Pk.*

N. Y. State Mus. Bul. 122, p.138, pl.110, fig.7-14. 1908

Russula roseipes (*Secr.*) *Bres.*

N. Y. State Mus. Rep't 51, p.306-7, pl.53, fig.1-7. 1898

N. Y. State Mus. Mem. 4, p.153-54, pl. 54, fig.1-7. 1900

Russula rugulosa *Pk.*

N. Y. State Mus. Rep't 54, p.179-80, pl.72, fig. 12-18. 1901

Russula sordida *Pk.*

N. Y. State Mus. Bul. 105, p.39-40, pl.98. 1906

Russula subsordida *Pk.*

N. Y. State Mus. Bul. 105, p.40-41, pl.99. 1906

Russula uncialis *Pk.*

N. Y. State Mus. Bul. 116, p.43, pl.107, fig.7-12. 1907

- Russula variata* *Banning*
N. Y. State Mus. Bul. 105, p.41-42, pl.101. 1906
- Russula virescens* *Fr.*
N. Y. State Mus. Rep't 48, p.189-90, pl.31. 1896. Bot. ed.
- Russula viridella* *Pk.*
N. Y. State Mus. Bul. 105, p.41, pl.100. 1906
- Strobilomyces strobilaceus* (*Scop.*) *Berk.*
N. Y. State Mus. Bul. 94, p.48-49, pl.92. 1905
- Stropharia bilamellata* *Pk.*
N. Y. State Mus. Bul. 122, p.139-40, pl.112, fig.5-10. 1908
- Tricholoma hirtellum* *Pk.*
N. Y. State Mus. Bul. 116, p.38-39, pl.105, fig.1-5. 1907
- Tricholoma imbricatum* *Fr.*
N. Y. State Mus. Rep't 48, p.169-70, pl.21, fig.6-11. 1896. Bot. ed.
- Tricholoma nudum* (*Bull.*) *Fr.*
N. Y. State Mus. Bull. 116, p.39-40, pl.104. 1907
- Tricholoma personatum* *Fr.*
N. Y. State Mus. Rep't 48, p.170-72, pl.22. 1896. Bot. ed.
- Tricholoma portentosum centrale* *Pk.*
N. Y. State Mus. Bul. 25, p.673, pl.57, fig.1-5. 1899
- Tricholoma portentosum centrale* *Pk.*
N. Y. State Mus. Mem. 4, p.138-39, pl.45, fig.1-5. 1900
- Tricholoma radiculatum* *Pk.*
N. Y. State Mus. Bul. 67, p.40-41, pl.82, fig.15-19. 1903
- Tricholoma russula* (*Schaeff.*) *Fr.*
N. Y. State Mus. Bul. 54, p.966-67, pl.77, fig.1-5. 1902
- Tricholoma silvaticum* *Pk.*
N. Y. State Mus. Bul. 67, p.41, pl.82, fig.1-6. 1903
- Tricholoma sordidum* (*Schum.*) *Fr.*
N. Y. State Mus. Bul. 131, p.38-39, pl.115. 1909
- Tricholoma subacutum* *Pk.*
N. Y. State Mus. Bul. 67, p.39-40, pl.82, fig.7-14. 1903
- Tricholoma terreum fragrans* *Pk.*
N. Y. State Mus. Rep't 49, p.57, pl.47, fig.1-10. 1896. Bot. ed.
- Tricholoma terreum fragrans* *Pk.*
N. Y. State Mus. Mem. 4, p.137-38, pl.45, fig.6-15. 1900
- Tricholoma transmutans* *Pk.*
N. Y. State Mus. Rep't 48, p.168-69, pl.21, fig.1-5. 1896. Bot. ed.
- Tricholoma unifactum* *Pk.*
N. Y. State Mus. Bul. 105, p.36-37, pl.94. 1906

Poisonous or unwholesome

- Amanita muscaria* *L.*
N. Y. State Mus. Rep't 48, p.212-15, pl.42. 1896. Bot. ed.
- Amanita phalloides* *Fr.*
N. Y. State Mus. Rep't 48, p.215-17, pl.40, 41, fig.1-3. 1896. Bot. ed.
- Amanita verna* *Bull.*
N. Y. State Mus. Bul. 48, p.215, pl.41, fig.4-7. 1896. Bot. ed.
- Boletus felleus* *Bull.*
N. Y. State Mus. Rep't 48, p.217-18, pl.43. 1896. Bot. ed.
- Clitocybe illudens* *Schw.*
N. Y. State Mus. Rep't 49, p.65, pl.49. 1896. Bot. ed.
- Clitocybe illudens* *Schw.*
N. Y. State Mus. Mem. 4, p.179-80, pl.68. 1900

LIST OF GENERA WHOSE NEW YORK SPECIES
(CHIEFLY) HAVE BEEN COLLATED WITH DESCRIPTIONS IN THE STATE BOTANIST'S REPORTS CITED

Aecidium

N. Y. State Mus. Rep't 24, p.105-8. 1872

Agaricus

N. Y. State Cab. Rep't 23, p.96-98. 1872. Bot. ed.

N. Y. State Mus. Rep't 36, p.41-49. 1884

Amanita

N. Y. State Cab. Rep't 23, p.68-70. 1872. Bot. ed.

N. Y. State Mus. Rep't 33, p. 38-49. 1880

Armillaria

N. Y. State Cab. Rep't 23, p.73. 1872. Bot. ed.

N. Y. State Mus. Rep't 43, p.40-44. 1890. Bot. ed.

N. Y. State Mus. Rep't 43, p.44-45. 1890. Bot. ed. (United States species)

Boletus

N. Y. State Cab. Rep't 23, p.127-33. 1872. Bot. ed.

N. Y. State Mus. Bul. 2, p.57-66. 1887

N. Y. State Mus. Bul. 8, p.80-157. 1889. (United States species)

Boletinus

N. Y. State Mus. Bul. 8, p.74-80. 1889. (United States species)

Cantharellus

N. Y. State Cab. Rep't 23, p.121-24. 1872. Bot. ed.

N. Y. State Mus. Bul. 2, p.34-43. 1887

Claudopus

N. Y. State Mus. Rep't 39, p.67-69. 1886

Clavaria

N. Y. State Mus. Rep't 24, p.104-5. 1872

Clitocybe

N. Y. State Cab. Rep't 23, p.75-78. 1872. Bot. ed.

Clitopilus

N. Y. State Mus. Rep't 42, p.39-46. 1889. Bot. ed.

Collybia

N. Y. State Cab. Rep't 23, p.78-80. 1872. Bot. ed.

N. Y. State Mus. Rep't 49, p. 32-55. 1896. Bot. ed.

Coprinus

N. Y. State Cab. Rep't 23, p.103-4. 1872. Bot. ed.

Cortinarius

N. Y. State Cab. Rep't 23, p.105-12. 1872. Bot. ed.

Craterellus

N. Y. State Mus. Bul. 2, p.44-48. 1887

Crepidotus

N. Y. State Mus. Rep't 39, p.69-73. 1886

Entoloma

N. Y. State Cab. Rep't 23, p.88. 1872. Bot. ed.

N. Y. State Mus. Bul. 131, p.47-54. 1909

Flammula

N. Y. State Cab. Rep't 23, p.90-91. 1872. Bot. ed.

N. Y. State Mus. Rep't 50, p.133-44. 1897

Galera

N. Y. State Cab. Rep't 23, p.93-94. 1872. Bot. ed.

N. Y. State Mus. Rep't 46, p.61-69. 1893. Bot. ed.

Hebeloma

N. Y. State Cab. Rep't 23, p.95-96. 1872. Bot. ed.

Helvella

N. Y. State Mus. Rep't, 31, p.60. 1879

Hygrophorus

N. Y. State Cab. Rep't 23, p.112-14. 1872. Bot. ed.

N. Y. State Mus. Bul. 116, p.45-67. 1907

Hypholoma

N. Y. State Cab. Rep't 23, p.98-99. 1872. Bot. ed.

Lactarius

N. Y. State Cab. Rep't 23, p.114-20. 1872. Bot. ed.

N. Y. State Mus. Rep't 38, p.111-33. 1885

Lentinus

N. Y. State Cab. Rep't 23, p.126-27. 1872. Bot. ed.

N. Y. State Mus. Bul. 131, p.42-47. 1909

Lepiota

N. Y. State Cab. Rep't 23, p.70-73. 1872. Bot. ed.

N. Y. State Mus. Rep't 35, p.150-64. 1884

Leptonia

N. Y. State Cab. Rep't 23, p.89. 1872. Bot. ed.

Lycoperdon

N. Y. State Mus. Rep't 32, p.58-72. 1879

Marasmius

N. Y. State Cab. Rep't 23, p.124-26. 1872. Bot. ed.

Mycena

N. Y. State Cab. Rep't 23, p.80-84. 1872. Bot. ed.

Naucoria

N. Y. State Cab. Rep't 23, p.91-93. 1872. Bot. ed.

Odontia

N. Y. State Mus. Rep't 53, p.847. 1900

Omphalia

N. Y. State Cab. Rep't 23, p.84-85. 1872. Bot. ed.

N. Y. State Mus. Rep't 45, p.32-42. 1893. Bot. ed.

Panaeolus

N. Y. State Cab. Rep't 23, p.100-2. 1872. Bot. ed.

Paxillus

N. Y. State Mus. Bul. 2, p.29-33. 1887

Pholiota

N. Y. State Cab. Rep't 23, p.89-90. 1872. Bot. ed.

N. Y. State Mus. Bul. 122, p.141-58. 1908

Pleurotus

N. Y. State Cab. Rep't 23, p.86-87. 1872. Bot. ed.

N. Y. State Mus. Rep't 39, p.58-67. 1886

Pluteolus

N. Y. State Mus. Rep't 46, p.58-61. 1893. Bot. ed.

Pluteus

N. Y. State Cab. Rep't 23, p.87-88. 1872. Bot. ed.

N. Y. State Mus. Rep't 38, p.133-38. 1885

Psathyrella

N. Y. State Cab. Rep't 23, p.102-3. 1872. Bot. ed.

Psilocybe

N. Y. State Cab. Rep't 23, p.99-100. 1872. Bot. ed.

Puccinia

N. Y. State Mus. Rep't 25, p.110-23. 1873

Russula

N. Y. State Cab. Rep't 23, p.120-21. 1872. Bot. ed.

N. Y. State Mus. Bul. 116, p.67-98. 1907

Spathularia

N. Y. State Mus. Rep't 50, p.118-19. 1897

Strobilomyces

N. Y. State Mus. Bul. 8, p.158-59. 1889. (United States species)

Trametes

N. Y. State Mus. Rep't 54, p.169-70. 1901

Tricholoma

N. Y. State Cab. Rep't 23, p.73-75. 1872. Bot. ed.

N. Y. State Mus. Rep't 44, p.38-64. 1891. Bot. ed.

Xylaria

N. Y. State Mus. Rep't 31, p.59. 1879

EXPLANATION OF PLATES

PLATE II

Hypholoma boughtoni Pk.

BOUGHTON HYPHOLOMA

- 1 Cluster of three immature plants
- 2 Mature umbonate plant
- 3 Mature plant without an umbo
- 4 Vertical section of upper part of an immature plant
- 5 Vertical section of upper part of a mature plant
- 6 Transverse section of a stem
- 7 Four spores, x 400



HYPHOLOMA BOUGHTONI Pk.
BOUGHTON HYPHOLOMA

PLATE III

Hypholoma rigidipes Pk.

RIGID STEM HYPHOLOMA

- 1 Immature plant
- 2 Mature plant
- 3 Vertical section of upper part of an immature plant
- 4 Vertical section of upper part of a mature plant
- 5 Transverse section of a stem
- 6 Four spores, x 400

Psilocybe nigrella Pk. •

BLACKISH PSILOCYBE

- 7 Immature plant with moist cap
- 8 Mature plant with moist cap
- 9 Mature plant with dry cap
- 10 Vertical section of upper part of a mature plant
- 11 Four spores, x 400



FIG. 1-6

HYPHOLOMA RIGIDIPES PK.
RIGID STEM *HYPHOLOMA*

FIG. 7-11

PSILOCYBE NIGRELLA PK.
BLACKISH *PSILOCYBE*

PLATE 117

Hebeloma album Pk.

WHITE HEBELOMA

- 1 Young plant
- 2 Mature plant with expanded cap
- 3 Mature plant with convex cap tinged with yellow
- 4 Vertical section of upper part of a young plant
- 5 Vertical section of upper part of a mature plant
- 6 Four spores, x 400

Clitocybe multiceps Pk.

MANY CAP CLITOCYBE

- 7 Cluster of seven plants
- 8 Vertical section of upper part of a plant
- 9 Four spores, x 400



FIG. 1-6

HEBELOMA ALBUM PK.
WHITE HEBELOMA

FIG. 7-9

CLITOCYBE MULTICEPS PK.
MANY CAP CLITOCYBE

PLATE 118

Lactarius aquifluus Pk.

WATERY MILK LACTARIUS

- 1 Young plant with moist cap
- 2 Mature plant with dry margin
- 3 Mature plant with entire cap dry
- 4 Vertical section of upper part of a plant
- 5 Transverse section of a stem
- 6 Four spores, x 400




LACTARIUS AQUIFLUUS Pk.
WATERY MILK LACTARIUS

PLATE 119

Entoloma grande Pk.

GRAND ENTOLOMA

- 1 Cluster of one mature and two young plants
- 2 Mature plant with umbonate and rugosely wrinkled cap .
- 3 Vertical section of upper part of a young plant
- 4 Vertical section of upper part of a mature plant
- 5 Four spores, x 400



ENTOLOMA GRANDE Pk.
GRAND ENTOLOMA

PLATE 120

Boletus viridarius Frost

GREEN LAWN BOLETUS

- 1 Young plant with tubes concealed by the veil
- 2 Young plant with tubes exposed
- 3 Mature plant showing a fragment of the white veil still adhering to the margin of the cap
- 4 Mature plant with whitish cap appendiculate with the ruptured veil
- 5 Young plant with yellowish tubes
- 6, 7 Mature plants showing color of mature tubes; stem of no. 7 only partly reticulated above the collar
- 8 Vertical section of upper part of a young plant
- 9 Vertical section of upper part of a mature plant
- 10 Four spores, x 400



BOLETUS VIRIDARIUS FROST
GREEN LAWN BOLETUS

EXTRALIMITAL SPECIES

PLATE W

Amanita morrisii Pk. .

MORRIS AMANITA

- 1 Young plant with expanded cap
- 2 Mature plant with expanded cap and two fragments of the volva adhering to the base of the stem
- 3 Vertical section of the upper part of a mature plant
- 4 Four spores, x 400



AMANITA MORRISII PK.
MORRIS AMANITA

PLATE X

Lactarius bryophilus Pk.

MOSS LOVING LACTARIUS

- 1 Young plant
- 2 Mature plant showing upper surface of cap
- 3 Vertical section of a young plant
- 4 Vertical section of a mature plant
- 5 Four spores, x 400

Agaricus eludens Pk.

ELUSIVE MUSHROOM

- 6 Young plant showing white gills and brown cap
- 7 Middle aged plant showing pink gills and brown cap
- 8 Mature plant showing brown gills and scaly cap
- 9 Mature plant showing red wound spot on the stem
- 10 Vertical section of upper part of a middle aged plant
- 11 Vertical section of upper part of a mature plant
- 12 Transverse section of a stem
- 13 Four spores, x 400



FIG. 1-5

LACTARIUS BRYOPHILUS Pk.
MOSS LOVING LACTARIUS

FIG. 6-13

AGARICUS ELUDENS Pk.
ELUDING AGARICUS

PLATE Y

Cortinarius ferrugineo-griseus Pk.

RUSTY GRAY CORTINARIUS

- 1 Young moist plant showing the webby veil
- 2 Mature moist plant showing remains of the spore stained veil on the stem
- 3 Vertical section of a young plant
- 4 Vertical section of a mature plant with part of the cap wanting



G. E. M. del.

CORTINARIUS FERRUGINEO-GRISEUS Pk.
RUSTY GRAY CORTINARIUS

PLATE Z

Cortinarius ferrugineo-griseus Pk.

RUSTY GRAY CORTINARIUS

- 1 Mature dry plant with violaceous stem
- 2 Vertical section of a small mature plant with violaceous stem
- 3 Four spores, x 400

Cortinarius actutoides Pk.

ACUTOID CORTINARIUS

- 4 Group of six young moist plants, one showing floccose scales of the white veil on the cap
- 5 Two mature dry plants
- 6 Vertical section of a young plant
- 7 Vertical section of a mature plant
- 8 Four spores, x 400

Russula blackfordae Pk.

BLACKFORD RUSSULA

- 9, 10 Two plants with convex cap
- 11 Mature plant with expanded cap
- 12 Vertical section of a mature plant
- 13 Four spores, x 400



FIG. 1-3

FIG. 4-8

CORTINARIUS FERRUGINEO-GRISEUS PK. CORTINARIUS ACUTOIDES PK.
RUSTY GRAY CORTINARIUS ACUTOID CORTINARIUS

FIG. 9-13

RUSSULA BLACKFORDAE PK.
BLACKFORD RUSSULA

INDEX

Agaricus campester, 7.
hortensis, 7.
eludens, 42.
explanation of plate, 106.

Agropyrum tenerum, 19.
Amanita morrisii, 42.
explanation of plate, 104.
Ascochyta solani-nigri, 19.

Belonidium glyceriae, 19.
Biatora cupreo-rosella, 19.
Bidens tenuisecta, 19.
Boletus viridarius, 19, 41.
explanation of plate, 102.
Bromus altissimus, 19.

Cardamine bulbosa, 33.
douglasii, 20.
Carduus crispus, 20.
Carex bebbii, 20.
crawfordii, 20.
Centaurea solstitialis, 33.
Cerastium viscosum, 33.
Chaenactis stevioides, 20.
Ciboria luteo-virescens, 20.
Clavaria lavendula, 47.
pallens, 47.
Clitocybe, many cap, 37-38.
Clitocybe candida, 20.
multiceps, 37.
explanation of plate, 96.
Cortinarius acutoides, 46.
explanation of plate, 110.
ferrugineo-griseus, 46.
explanation of plate, 108, 110.
subsalmoneus, 20.
Crataegus brevipes, 20.
verecunda gonocarpa, 33.

Denudata, 71.
Diplocladium penicilloides, 21.
Diplodia cercidis, 22.
hamamelidis, 22.
tamariscina, 22.

Discina leucoxantha, 22.
Dothiorella divergens, 22.

Edible fungi, 6, 37-41; list, 78-89.
Entoloma grande, 39.
explanation of plate, 100.
Epilobium densum, 34.
Epipactis tesselata, 23.
Erythronium albidum, 34.
Exoascus pruni, 34.
Explanation of plates, 91-110.

Fenestella amorphia, 23.
Flammula highlandensis, 77.
Fungi, edible, 6, 37-41; extralimital,
new species, 42-48; list, 78-89.

Galium erectum, 34.
Geum flavum, 23.

Hebeloma, New York species, 67-
77.
colvin, 73.
crustuline, 73.
disk diseased, 75.
firm, 70.
fragile, 77.
glutinous, 68.
gregarious, 75.
long stem, 73.
marsh, 76.
ochery, 70.
pale margin, 76.
pasture, 71.
pink gill, 71.
slightly sordid, 75.
small fruit, 70.
social, 74.
thin margin, 77.
unlawful, 76.
veiled, 69.
white, 40, 72.
whitish, 72.
Hebeloma, 67.
albidulum, 72.

Hebeloma (continued)

- album, 40, 72.
- explanation of plate, 96.
- colvini, 73.
- crustuliniforme, 73.
- discomorbidum, 75.
- excedens, 77.
- fastibile, 70.
- firmum, 70.
- fragilius, 77.
- glutinosum, 68.
- gregarium, 75.
- illicitum, 76.
- longicaudum, 73.
- pallidomarginatum, 76.
- palustre, 76.
- parvifructum, 70.
- pascuense, 71.
- sarcophyllum, 71.
- sociale, 74.
- sordidulum, 75.
- velatum, 69.
- Hedeoma hispida*, 34.
- Hypholoma boughtoni*, 23.
- explanation of plate, 92.
- rigidipes, 24.
- explanation of plate, 94.

Ilex verticillata tenuifolia, 34.*Indusiata*, 68.*Inocybe*, New York species, 48-67.

- agglutinate, 62.
- black disk, 64.
- brown disk, 66.
- changed, 50.
- chestnut, 58.
- cracked, 56.
- curved scale, 49.
- earthy leaf, 61.
- eutheloid, 57.
- excoriate, 55.
- fallacious, 63.
- feeble, 64.
- fibrillose, 50.
- grayish, 57.
- hairy cap, 60.
- hairy margin, 65.
- late, 61.
- mammillate, 56.
- maritimoid, 53.

Inocybe (continued)

- marsh, 63.
- ochraceous, 62.
- one colored, 50.
- pale stem, 55.
- rigid stem, 59.
- rough spore, 66.
- scaly disk, 53.
- small, 53.
- star spore, 59.
- stellate spore, 51.
- subtomentose, 62.
- tawny, 54.
- umbo marked, 58.
- unfortunate, 52.
- untrusty, 64.
- vatricosoid, 67.
- violaceous gill, 57.
- white disk, 59.
- woolly, 51.
- Inocybe*, 48.
- agglutinata, 62.
- albodisca, 59.
- asterospora, 59, 77.
- calamistrata, 49.
- castanea, 58.
- comatella, 60.
- diminuta, 53.
- eutheles, 56.
- eutheloides, 57.
- excoriata, 55.
- fallax, 63.
- fibrillosa, 50.
- fuscodisca, 66.
- geophylla, 61.
- griseoscabrosa, 57.
- infelix, 52, 77.
- var. brevipes*, 52.
- infida, 64.
- lanuginosa, 51.
- maritimoides, 53.
- mutata, 50.
- nigridisca, 64.
- pallidipes, 55.
- paludinella, 63.
- rigidipes, 59.
- rimosa, 56.
- var. parva*, 56.
- var. cuspidata*, 56.

Inocybe (*continued*)

- serotina, 61.
- squamosodisca, 53.
- stellatospora, 51.
- subexilis, 64.
- subfulva, 54.
- subochracea, 62.
- subtomentosa, 62.
- trechispora, 66.
- trichcloma, 65.
- umboninota, 58.
- unicolor, 50.
- vaticosoides, 67.
- violaceifolia, 57.

Juncus *brachycephalus*, 24.

- brevicaudatus*, 24.
- secundus*, 24.

Juniperus *horizontalis*, 24.**Lacerae**, 52.**Lactarius** *aquifluus*, 38.

- explanation of plate, 98.
- bryophilus*, 44.

- explanation of plate, 106.

Lactuca *scariola integrata*, 34.**Laportea** *canadensis*, 35.**Leontodon** *nudicaulis*, 25.**Leskea** *gracilescens*, 25.**Ligusticum** *scoticum*, 25.**Listera** *australis*, 35.**Lophiotrema** *hystericides*, 25.

- littorale*, 25.

Marasmius *alienus*, 25.

- oreades*, 35.

Melanopsamma *confertissima*, 25.**Micocera** *coccophila*, 25.**Midotis** *irregularis*, 26.**Monolepis** *nuttalliana*, 26.**Morchella** *crispa*, 26.

- rimosipes*, 26.

Mushrooms, *see* **Fungi**.**Naias** *gracillima*, 26.**Nardia** *crenulata*, 26.

- hyalira*, 26.

Naucoria *sphagnophila*, 45.**Omphala** *rugosodisca levidisca*, 35.**Panicum** *implicatum*, 26.

- oricola*, 26.

- spretum*, 27.

Peridermium *consimile*, 35.

- strobi*, 27.

Pezizella *lanceolato-paraphysata*, 27.**Phaeopezia** *fuscocarpa*, 27.**Pholiota** *aurivella*, 27.**Phomopsis** *stewartii*, 27.**Picris** *echinoides*, 28.**Plants**, species added to collection,

- 5, 6, 8-10; species not before reported, 5, 19-32; contributors and their contributions, 10-18.

Plates, explanation, 91-110.**Polyporus** *giganteus*, 35.**Potamogeton** *richardsoni*, 28.**Prunus** *pumila*, 35.**Psilocybe** *nigrella*, 28.

- explanation of plate, 94.

Puccinia *epiphylla*, 28.**Pusilla**, 74.**Pyrus** *coronaria*, 36.

- melanocarpa*, 36.

Ribes *triste albinervium*, 28.**Rimosae**, 54.**Rubia** *tinctorum*, 29.**Rubus** *andrewsianus*, 29.

- permixtus*, 29.

- recurvans*, 29.

Rumex *pallidus*, 29.**Russula** *blackfordae*, 43.

- explanation of plate, 110.

- serissima*, 44.

Schwalbea *americana*, 36.**Septoria** *sedicola*, 29.**Solanum** *nigrum*, 36.**Solidago** *aspera*, 30.

- squarrosa ramosa*, 36.

Sparganium *americanum*, 30.

- angustifolium*, 30.

- diversifolium*, 30.

Squarrosae, 49.**Stachys** *sieboldii*, 30.**Stephanoma** *strigosum*, 30.**Sterigmatocystis** *ochracea*, 30.

Thalictrum confine, 37.

revolutum, 37.

Trametes merisma, 31.

Trichosporium variabile, 31.

Velutinae, 60.

Verticillium rexianum, 32.

Viola sororia, 37.

Viscidae, 65.

Volvaria volvacea, 32.

Zizania palustris, 32.

New York State Education Department

New York State Museum

JOHN M. CLARKE, Director

PUBLICATIONS

Packages will be sent prepaid except when distance or weight renders the same impracticable. On 10 or more copies of any one publication 20% discount will be given. Editions printed are only large enough to meet special claims and probable sales. When the sale copies are exhausted, the price for the few reserve copies is advanced to that charged by second-hand booksellers, in order to limit their distribution to cases of special need. Such prices are inclosed in []. All publications are in paper covers, unless binding is specified. Checks or money orders should be addressed and payable to New York State Education Department.

Museum annual reports 1847-date. *All in print to 1894, 50c a volume, 75c in cloth; 1894-date, sold in sets only; 75c each for octavo volumes; price of quarto volumes on application.*

These reports are made up of the reports of the Director, Geologist, Paleontologist, Botanist and Entomologist, and museum bulletins and memoirs, issued as advance sections of the reports.

Director's annual reports 1904-date.

1904. 138p. 20c.

1905. 102p. 23pl. 30c.

1906. 186p. 41pl. 35c.

1907. 212p. 63pl. 50c.

1908. 234p. 39pl. map. 40c.

These reports cover the reports of the State Geologist and of the State Paleontologist. Bound also with the museum reports of which they form a part.

Geologist's annual reports 1881-date. Rep'ts 1, 3-13, 17-date, 8vo; 2, 14-16, 4to.

In 1898 the paleontologic work of the State was made distinct from the geologic and was reported separately from 1899-1903. The two departments were reunited in 1904, and are now reported in the Director's report.

The annual reports of the original Natural History Survey, 1837-41, are out of print.

Reports 1-4, 1881-84, were published only in separate form. Of the 5th report 4 pages were reprinted in the 39th museum report, and a supplement to the 6th report was included in the 40th museum report. The 7th and subsequent reports are included in the 41st and following museum reports, except that certain lithographic plates in the 11th report (1891) and 13th (1893) are omitted from the 45th and 47th museum reports.

Separate volumes of the following only are available.

Report	Price	Report	Price	Report	Price
12 (1892)	\$.50	17	\$.75	21	\$.40
14	.75	18	.75	22	.40
15, 2v.	2	19	.40	23	.45
16	1	20	.50		

[See Director's annual reports]

Paleontologist's annual reports 1899-date.

See first note under Geologist's annual reports.

Bound also with museum reports of which they form a part. Reports for 1899 and 1900 may be had for 20c each. Those for 1901-3 were issued as bulletins. In 1904 combined with the Director's report.

Entomologist's annual reports on the injurious and other insects of the State of New York 1882-date.

Reports 3-20 bound also with museum reports 40-46, 48-58 of which they form a part. Since 1898 these reports have been issued as bulletins. Reports 3-4, 17 are out of print, other reports with prices are:

Report	Price	Report	Price	Report	Price
1	\$.50	10	\$.35	18 (Bul. 64)	\$.20
2	.30	11	.25	19 (" 76)	.15
5	.25	12	.25	20 (" 97)	.40
6	.15	13	Free	21 (" 104)	.25
7	.20	14 (Bul. 23)	.20	22 (" 110)	.25
8	.25	15 (" 31)	.15	23 (" 124)	.75
9	.25	16 (" 36)	.25	24 (" 134)	.35

Reports 2, 8-12 may also be obtained bound in cloth at 25c each in addition to the price given above.

Botanist's annual reports 1867-date.

Bound also with museum reports 21-date of which they form a part; the first Botanist's report appeared in the 21st museum report and is numbered 21. Reports 21-24, 29, 31-41 were not published separately.

Separate reports for 1871-74, 1876, 1888-98 are out of print. Report for 1899 may be had for 20c; 1900 for 50c. Since 1901 these reports have been issued as bulletins.

NEW YORK STATE EDUCATION DEPARTMENT

Descriptions and illustrations of edible, poisonous and unwholesome fungi of New York have also been published in volumes 1 and 3 of the 48th (1894) museum report and in volume 1 of the 49th (1895), 51st (1897), 52d (1898), 54th (1900), 55th (1901), in volume 4 of the 56th (1902), in volume 2 of the 57th (1903), in volume 4 of the 58th (1904), in volume 2 of the 59th (1905), 60th (1906), in volume 2 of the 61st (1907) and 62d (1908) reports. The descriptions and illustrations of edible and unwholesome species contained in the 49th, 51st and 52d reports have been revised and rearranged, and, combined with others more recently prepared, constitute Museum memoir 4.

Museum bulletins 1887-date. 8vo. *To advance subscribers, \$2 a year or \$1 a year for division (1) geology, economic geology, paleontology, mineralogy; 50c each for divisions (2) general zoology, archeology and miscellaneous, (3) botany, (4) entomology.*

Bulletins are grouped in the list on the following pages according to divisions.

The divisions to which bulletins belong are as follows:

1 Zoology	48 Geology	94 Botany
2 Botany	49 Paleontology	95 Geology
3 Economic Geology	50 Archeology	96 " "
4 Mineralogy	51 Zoology	97 Entomology
5 Entomology	52 Paleontology	98 Mineralogy
6 " "	53 Entomology	99 Paleontology
7 Economic Geology	54 Botany	100 Economic Geology
8 Botany	55 Archeology	101 Paleontology
9 Zoology	56 Geology	102 Economic Geology
10 Economic Geology	57 Entomology	103 Entomology
11 " "	58 Mineralogy	104 " "
12 " "	59 Entomology	105 Botany
13 Entomology	60 Zoology	106 Geology
14 Geology	61 Economic Geology	107 " "
15 Economic Geology	62 Miscellaneous	108 Archeology
16 Archeology	63 Paleontology	109 Entomology
17 Economic Geology	64 Entomology	110 " "
18 Archeology	65 Paleontology	111 Geology
19 Geology	66 Miscellaneous	112 Economic Geology
20 Entomology	67 Botany	113 Archeology
21 Geology	68 Entomology	114 Paleontology
22 Archeology	69 Paleontology	115 Geology
23 Entomology	70 Mineralogy	116 Botany
24 " "	71 Zoology	117 Archeology
25 Botany	72 Entomology	118 Paleontology
26 Entomology	73 Archeology	119 Economic Geology
27 " "	74 Entomology	120 " "
28 Botany	75 Botany	121 Director's report for 1907
29 Zoology	76 Entomology	122 Botany
30 Economic Geology	77 Geology	123 Economic Geology
31 Entomology	78 Archeology	124 Entomology
32 Archeology	79 Entomology	125 Archeology
33 Zoology	80 Paleontology	126 Geology
34 Paleontology	81 " "	127 " "
35 Economic Geology	82 " "	128 Paleontology
36 Entomology	83 Geology	129 Entomology
37 " "	84 " "	130 Zoology
38 Zoology	85 Economic Geology	131 Botany
39 Paleontology	86 Entomology	132 Economic Geology
40 Zoology	87 Archeology	133 Director's report for 1908
41 Archeology	88 Zoology	134 Entomology
42 Paleontology	89 Archeology	135 Geology
43 Zoology	90 Paleontology	136 Entomology
44 Economic Geology	91 Zoology	137 Geology
45 Paleontology	92 Paleontology	138 " "
46 Entomology	93 Economic Geology	139 Botany
47 " "		

Bulletins are also found with the annual reports of the museum as follows:

Bulletin	Report	Bulletin	Report	Bulletin	Report	Bulletin	Report
12-15	48, v. 1	69	56, v. 2	97	58, v. 5	125	62, v. 3
16, 17	50, v. 1	70, 71	57, v. 1, pt 1	98, 99	59, v. 2	126-28	62, v. 1
18, 19	51, v. 1	72	57, v. 1, pt 2	100	59, v. 1	129	62, v. 2
20-25	52, v. 1	73	57, v. 2	101	59, v. 2	130	62, v. 3
26-31	53, v. 1	74	57, v. 1, pt 2	102	59, v. 1	131, 132	62, v. 2
32-34	54, v. 1	75	57, v. 2	103-5	59, v. 2	133	62, v. 1
35, 36	54, v. 2	76	57, v. 1, pt 2	106	59, v. 1	134	62, v. 2
37-44	54, v. 3	77	57, v. 1, pt 1	107	60, v. 2		
45-48	54, v. 4	78	57, v. 2	108	60, v. 3		
49-54	55, v. 1	79	57, v. 1, pt 2	109, 110	60, v. 1		
55	56, v. 4	80	57, v. 1, pt 1	111	60, v. 2		
56	56, v. 1	81, 82	58, v. 3	112	60, v. 1	<i>Memoir</i>	
57	56, v. 3	83, 84	58, v. 1	113	60, v. 3		
58	56, v. 1	85	58, v. 2	114	60, v. 1	2	49, v. 3
59, 60	56, v. 3	86	58, v. 5	115	60, v. 2	3, 4	53, v. 2
61	56, v. 1	87-89	58, v. 4	116	60, v. 1	5, 6	57, v. 3
62	56, v. 4	90	58, v. 3	117	60, v. 3	7	57, v. 4
63	56, v. 2	91	58, v. 4	118	60, v. 1	8, pt 1	59, v. 3
64	56, v. 3	92	58, v. 3	119-21	61, v. 1	8, pt 2	59, v. 4
65	56, v. 2	93	58, v. 2	122	61, v. 2	9, pt 1	60, v. 4
66, 67	56, v. 4	94	58, v. 4	123	61, v. 1	9, pt 2	62, v. 4
68	56, v. 3	95, 96	58, v. 1	124	61, v. 2	10	60, v. 5
						11	61, v. 3

MUSEUM PUBLICATIONS

The figures at the beginning of each entry in the following list indicate its number as a museum bulletin.

- Geology.** 14 Kemp, J. F. Geology of Moriah and Westport Townships, Essex Co. N. Y., with notes on the iron mines. 38p. il. 7pl. 2 maps. Sept. 1895. Free.
- 19 Merrill, F. J. H. Guide to the Study of the Geological Collections of the New York State Museum. 164p. 119pl. map. Nov. 1898. *Out of print.*
- 21 Kemp, J. F. Geology of the Lake Placid Region. 24p. 1pl. map. Sept. 1898. Free.
- 48 Woodworth, J. B. Pleistocene Geology of Nassau County and Borough of Queens. 58p. il. 8pl. map. Dec. 1901. 25c.
- 56 Merrill, F. J. H. Description of the State Geologic Map of 1901. 42p. 2 maps, tab. Nov. 1902. Free.
- 77 Cushing, H. P. Geology of the Vicinity of Little Falls, Herkimer Co. 98p. il. 15pl. 2 maps. Jan. 1905. 30c.
- 83 Woodworth, J. B. Pleistocene Geology of the Mooers Quadrangle. 62p. 25pl. map. June 1905. 25c.
- 84 ——— Ancient Water Levels of the Champlain and Hudson Valleys. 206p. il. 11pl. 18 maps. July 1905. 45c.
- 95 Cushing, H. P. Geology of the Northern Adirondack Region. 188p. 15pl. 3 maps. Sept. 1905. 30c.
- 96 Ogilvie, I. H. Geology of the Paradox Lake Quadrangle. 54p. il. 17pl. map. Dec. 1905. 30c.
- 106 Fairchild, H. L. Glacial Waters in the Erie Basin. 88p. 14pl. 9 maps. Feb. 1907. *Out of print.*
- 107 Woodworth, J. B.; Hartnagel, C. A.; Whitlock, H. P.; Hudson, G. H.; Clarke, J. M.; White, David & Berkey, C. P. Geological Papers. 388p. 54pl. map. May 1907. 90c, *cloth.*
- Contents:* Woodworth, J. B. Postglacial Faults of Eastern New York.
Hartnagel, C. A. Stratigraphic Relations of the Oneida Conglomerate.
——— Upper Siluric and Lower Devonian Formations of the Skunkmunk Mountain Region.
Whitlock, H. P. Minerals from Lyon Mountain, Clinton Co.
Hudson, G. H. On Some Pelmatozoa from the Chazy Limestone of New York.
Clarke, J. M. Some New Devonian Fossils.
——— An Interesting Style of Sand-filled Vein.
——— Eurypterid Shales of the Shawangunk Mountains in Eastern New York.
White, David. A Remarkable Fossil Tree Trunk from the Middle Devonian of New York.
Berkey, C. P. Structural and Stratigraphic Features of the Basal Gneisses of the Highlands.
- 111 Fairchild, H. L. Drumlins of New York. 60p. 28pl. 19 maps. July 1907. *Out of print.*
- 115 Cushing, H. P. Geology of the Long Lake Quadrangle. 88p. 20pl. map. Sept. 1907. 25c.
- 126 Miller, W. J. Geology of the Remsen Quadrangle. 54p. il. 11pl. map. Jan. 1909. 25c.
- 127 Fairchild, H. L. Glacial Waters in Central New York. 64p. 27pl. 15 maps. Mar. 1909. 40c.
- 135 Miller, W. J. Geology of the Port Leyden Quadrangle, Lewis County, N. Y., 62p. il. 11 pl. map. Jan. 1910. 25c.
- 137 Luther, D. D. Geology of the Auburn-Genoa Quadrangles. 36p. map. Mar. 1910. 20c.
- 138 Kemp, J. F. & Ruedemann, Rudolf. Geology of the Elizabethtown and Port Henry Quadrangles. 176p. il. 20 pl. 3 maps. Apr. 1910. 40c.
- Berkey, C. P. Geology of the Highlands of the Hudson. *In preparation.*
- Cushing, H. P.; Fairchild, H. L.; Ruedemann, Rudolf & Smyth, C. H. Geology of the Thousand Island Region. *In press.*
- Economic geology.** 3 Smock, J. C. Building Stone in the State of New York. 154p. Mar. 1888. *Out of print.*
- 7 ——— First Report on the Iron Mines and Iron Ore Districts in the State of New York. 78p. map. June 1889. *Out of print.*
- 10 ——— Building Stone in New York. 210p. map, tab. Sept. 1890. 40c.
- 11 Merrill, F. J. H. Salt and Gypsum Industries of New York. 94p. 12pl. 2 maps, 11 tab. Apr. 1893. [50c]
- 12 Ries, Heinrich. Clay Industries of New York. 174p. 1pl. il. map. Mar. 1895. 30c.

NEW YORK STATE EDUCATION DEPARTMENT

- 15 Merrill, F. J. H. Mineral Resources of New York. 240p. 2 maps. Sept. 1895. [50c]
- 17 ——— Road Materials and Road Building in New York. 52p. 14pl. 2 maps. Oct. 1897. 15c.
- 30 Orton, Edward. Petroleum and Natural Gas in New York. 136p. il. 3 maps. Nov. 1899. 15c.
- 35 Ries, Heinrich. Clays of New York; their Properties and Uses 456p. 140pl. map. June 1900. *Out of print.*
- 44 ——— Lime and Cement Industries of New York; Eckel, E. C. Chapters on the Cement Industry. 332p. 101pl. 2 maps. Dec. 1901. 85c, *cloth.*
- 61 Dickinson, H. T. Quarries of Bluestone and other Sandstones in New York. 114p. 18pl. 2 maps. Mar. 1903. 35c.
- 85 Rafter, G. W. Hydrology of New York State. 902p. il. 44pl. 5 maps. May 1905. \$1.50, *cloth.*
- 93 Newland, D. H. Mining and Quarry Industry of New York. 78p. July 1905. *Out of print.*
- 100 McCourt, W. E. Fire Tests of Some New York Building Stones. 40p. 26pl. Feb 1906. 15c.
- 102 Newland, D. H. Mining and Quarry Industry of New York 1905. 162p. June 1906. *Out of print.*
- 112 ——— Mining and Quarry Industry of New York 1906. 82p. July 1907. *Out of print.*
- 119 ——— & Kemp, J. F. Geology of the Adirondack Magnetic Iron Ores with a Report on the Mineville-Port Henry Mine Group. 184p. 14pl. 8 maps. Apr. 1908. 35c.
- 120 Newland, D. H. Mining and Quarry Industry of New York 1907. 82p. July 1908. 15c.
- 123 ——— & Hartnagel, C. A. Iron Ores of the Clinton Formation in New York State. 76p. il. 14 pl. 3 maps. Nov. 1908. 25c.
- 132 Newland, D. H. Mining and Quarry Industry of New York 1908. 98p. July 1909. 15c.
- Mining and Quarry Industry of New York for 1909. *In preparation.*
- Mineralogy. 4 Nason, F. L. Some New York Minerals and their Localities. 22p. 1pl. Aug. 1888. Free.
- 58 Whitlock, H. P. Guide to the Mineralogic Collections of the New York State Museum. 150p. il. 39pl. 11 models. Sept. 1902. 40c.
- 70 ——— New York Mineral Localities. 110p. Oct. 1903. 20c.
- 98 ——— Contributions from the Mineralogic Laboratory. 38p. 7pl. Dec. 1905. 15c.
- Paleontology. 34 Cumings, E. R. Lower Silurian System of Eastern Montgomery County; Prosser, C. S. Notes on the Stratigraphy of Mohawk Valley and Saratoga County, N. Y. 74p. 14pl. map. May 1900. 15c.
- 39 Clarke, J. M.; Simpson, G. B. & Loomis, F. B. Paleontologic Papers 1. 72p. il. 16pl. Oct. 1900. 15c.
- Contents:* Clarke, J. M. A Remarkable Occurrence of Orthoceras in the Oneonta Beds of the Chenango Valley, N. Y.
—— Paropsonema cryptophya; a Peculiar Echinoderm from the Intumescens-zone (Portage Beds) of Western New York.
—— Dictyonine Hexactinellid Sponges from the Upper Devonian of New York.
—— The Water Biscuit of Squaw Island, Canandaigua Lake, N. Y.
Simpson, G. B. Preliminary Descriptions of New Genera of Paleozoic Rugose Corals.
Loomis, F. B. Siluric Fungi from Western New York.
- 42 Ruedemann, Rudolf. Hudson River Beds near Albany and their Taxonomic Equivalents. 116p. 2pl. map. Apr. 1901. 25c.
- 45 Grabau, A. W. Geology and Paleontology of Niagara Falls and Vicinity. 286p. il. 18pl. map. Apr. 1901. 65c; *cloth*, 90c.
- 49 Ruedemann, Rudolf; Clarke, J. M. & Wood, Elvira. Paleontologic Papers 2. 240p. 13pl. Dec. 1901. 40c.
- Contents:* Ruedemann, Rudolf. Trenton Conglomerate of Rysedorph Hill.
Clarke, J. M. Limestones of Central and Western New York Interbedded with Bituminous Shales of the Marcellus Stage.
Wood, Elvira. Marcellus Limestones of Lancaster, Erie Co., N. Y.
Clarke, J. M. New Agelacrinites.
—— Value of Amnigenia as an Indicator of Fresh-water Deposits during the Devonian of New York, Ireland and the Rhineland.

MUSEUM PUBLICATIONS

- 52 Clarke, J. M. Report of the State Paleontologist 1901. 28op. il. 1op. l. map, 1 tab. July 1902. 40c.
- 63 — & Luther, D. D. Stratigraphy of Canandaigua and Naples Quadrangles. 78p. map. June 1904. 25c.
- 65 Clarke, J. M. Catalogue of Type Specimens of Paleozoic Fossils in the New York State Museum. 848p. May 1903. \$1.20, cloth.
- 69 — Report of the State Paleontologist 1902. 464p. 52pl. 7 maps. Nov. 1903. \$1, cloth.
- 80 — Report of the State Paleontologist 1903. 396p. 29pl. 2 maps. Feb. 1905. 85c, cloth.
- 81 — & Luther, D. D. Watkins and Elmira Quadrangles. 32p. map. Mar. 1905. 25c.
- 82 — Geologic Map of the Tully Quadrangle. 40p. map. Apr. 1905. 20c.
- 90 Ruedemann, Rudolf. Cephalopoda of Beekmantown and Chazy Formations of Champlain Basin. 224p. il. 38pl. May 1906. 75c, cloth.
- 92 Grabau, A. W. Guide to the Geology and Paleontology of the Schoharie Region. 314p. il. 26pl. map. Apr. 1906. 75c, cloth.
- 99 Luther, D. D. Geology of the Buffalo Quadrangle. 32p. map. May 1906. 20c.
- 101 — Geology of the Penn Yan-Hammondsport Quadrangles. 28p. map. July 1906. 25c.
- 114 Hartnagel, C. A. Geologic Map of the Rochester and Ontario Beach Quadrangles. 36p. map. Aug. 1907. 20c.
- 118 Clarke, J. M. & Luther, D. D. Geologic Maps and Descriptions of the Portage and Nunda Quadrangles including a map of Letchworth Park. 50p. 16pl. 4 maps. Jan. 1908. 35c.
- 128 Luther, D. D. Geology of the Geneva-Ovid Quadrangles. 44p. map. Apr. 1909. 20c.
- Geology of the Phelps Quadrangle. *In preparation.*
- Whitnall, H. O. Geology of the Morrisville Quadrangle. *Prepared.*
- Hopkins, T. C. Geology of the Syracuse Quadrangle. *In preparation.*
- Hudson, G. H. Geology of Valcour Island. *In preparation.*
- Zoology. I Marshall, W. B. Preliminary List of New York Unionidae. 20p. Mar. 1892. Free.
- 9 — Beaks of Unionidae Inhabiting the Vicinity of Albany, N. Y. 30p. 1pl. Aug. 1890. Free.
- 20 Miller, G. S. jr. Preliminary List of New York Mammals. 124p. Oct. 1899. 15c.
- 33 Farr, M. S. Check List of New York Birds. 224p. Apr. 1900. 25c.
- 38 Miller, G. S. jr. Key to the Land Mammals of Northeastern North America. 106p. Oct. 1900. 15c.
- 40 Simpson, G. B. Anatomy and Physiology of Polygyra albolabris and Limax maximus and Embryology of Limax maximus. 82p. 28pl. Oct. 1901. 25c.
- 43 Kellogg, J. L. Clam and Scallop Industries of New York. 36p. 2pl. map. Apr. 1901. Free.
- 51 Eckel, E. C. & Paulmier, F. C. Catalogue of Reptiles and Batrachians of New York. 64p. il. 1pl. Apr. 1902. 15c.
- Eckel, E. C. Serpents of Northeastern United States.
- Paulmier, F. C. Lizards, Tortoises and Batrachians of New York.
- 60 Bean, T. H. Catalogue of the Fishes of New York. 784p. Feb. 1903. \$1, cloth.
- 71 Kellogg J. L. Feeding Habits and Growth of Venus mercenaria. 30p. 4pl. Sept. 1903. Free.
- 88 Letson, Elizabeth J. Check List of the Mollusca of New York. 116p. May 1905. 20c.
- 91 Paulmier, F. C. Higher Crustacea of New York City. 78p. il. June 1905. 20c.
- 130 Shufeldt, R. W. Osteology of Birds. 382p. il. 26pl. May 1909. 50c.
- Entomology. 5 Lintner, J. A. White Grub of the May Beetle. 34p. il. Nov. 1888. Free.
- 6 — Cut-worms. 38p. il. Nov. 1888. Free.

NEW YORK STATE EDUCATION DEPARTMENT

- 13 — San José Scale and Some Destructive Insects of New York State
54p. 7pl. Apr. 1895. 15c.
- 20 Felt, E. P. Elm-leaf Beetle in New York State. 46p. il. 5pl. June
1898. Free.
See 57.
- 23 — 14th Report of the State Entomologist 1898. 150p. il. 9pl. Dec.
1898. 20c.
- 24 — Memorial of the Life and Entomologic Work of J. A. Lintner Ph.D.
State Entomologist 1874-98; Index to Entomologist's Reports 1-13.
316p. 1pl. Oct. 1899. 35c.
Supplement to 14th report of the State Entomologist.
- 26 — Collection, Preservation and Distribution of New York Insects.
36p. il. Apr. 1899. Free.
- 27 — Shade Tree Pests in New York State. 26p. il. 5pl. May 1899.
Free.
- 31 — 15th Report of the State Entomologist 1899. 128p. June 1900.
15c.
- 36 — 16th Report of the State Entomologist 1900. 118p. 16pl. Mar.
1901. 25c.
- 37 — Catalogue of Some of the More Important Injurious and Beneficial
Insects of New York State. 54p. il. Sept. 1900. Free.
- 46 — Scale Insects of Importance and a List of the Species in New
York State. 94p. il. 15pl. June 1901. 25c.
- 47 Needham, J. G. & Betten, Cornelius. Aquatic Insects in the Adiron-
dacks. 234p. il. 36pl. Sept. 1901. 45c.
- 53 Felt, E. P. 17th Report of the State Entomologist 1901. 232p. il. 6pl.
Aug. 1902. *Out of print.*
- 57 — Elm Leaf Beetle in New York State. 46p. il. 8pl. Aug. 1902.
Out of print.
This is a revision of 20 containing the more essential facts observed since that was pre-
pared.
- 59 — Grapevine Root Worm. 40p. 6pl. Dec. 1902. 15c.
See 72.
- 64 — 18th Report of the State Entomologist 1902. 110p. 6pl. May
1903. 20c.
- 68 Needham, J. G. & others. Aquatic Insects in New York. 322p. 52pl.
Aug. 1903. 80c. *cloth.*
- 72 Felt, E. P. Grapevine Root Worm. 58p. 13pl. Nov. 1903. 20c.
This is a revision of 59 containing the more essential facts observed since that was
prepared.
- 74 — & Joutel, L. H. Monograph of the Genus Saperda. 88p. 14pl
June 1904. 25c.
- 76 Felt, E. P. 19th Report of the State Entomologist 1903. 150p. 4pl.
1904. 15c.
- 79 — Mosquitos or Culicidae of New York. 164p. il. 57pl. tab. Oct.
1904. 40c.
- 86 Needham, J. G. & others. May Flies and Midges of New York. 352p.
il. 37pl. June 1905. 80c. *cloth.*
- 97 Felt, E. P. 20th Report of the State Entomologist 1904. 246p. il. 19pl.
Nov. 1905. 40c.
- 103 — Gipsy and Brown Tail Moths. 44p. 10pl. July 1906. 15c.
- 104 — 21st Report of the State Entomologist 1905. 144p. 10pl. Aug.
1906. 25c.
- 109 — Tussock Moth and Elm Leaf Beetle. 34p. 8pl. Mar. 1907. 20c.
- 110 — 22d Report of the State Entomologist 1906. 152p. 3pl. June
1907. 25c.
- 124 — 23d Report of the State Entomologist 1907. 542p. 44pl. il.
Oct. 1908. 75c.
- 129 — Control of Household Insects. 48p. il. May 1909. *Out of print.*

- 134 — 24th Report of the State Entomologist 1908. 208p. 17pl. il. Sept. 1909. 35c.
- 136 — Control of Flies and Other Household Insects. 56p. il. Feb. 1910. 15c.
- This is a revision of 129 containing the more essential facts observed since that was prepared.
- Needham, J. G. Monograph on Stone Flies. *In preparation.*
- Botany. 2 Peck, C. H. Contributions to the Botany of the State of New York. 72p. 2pl. May 1887. *Out of print.*
- 8 — Boleti of the United States. 98p. Sept. 1889. *Out of print.*
- 25 — Report of the State Botanist 1898. 76p. 5pl. Oct. 1899. *Out of print.*
- 28 — Plants of North Elba. 206p. map. June 1899. 20c.
- 54 — Report of the State Botanist 1901. 58p. 7pl. Nov. 1902. 40c.
- 67 — Report of the State Botanist 1902. 196p. 5pl. May 1903. 50c.
- 75 — Report of the State Botanist 1903. 70p. 4pl. 1904. 40c.
- 94 — Report of the State Botanist 1904. 60p. 10pl. July 1905. 40c.
- 105 — Report of the State Botanist 1905. 108p. 12pl. Aug. 1906. 50c.
- 116 — Report of the State Botanist 1906. 120p. 6pl. July 1907. 35c.
- 122 — Report of the State Botanist 1907. 178p. 5pl. Aug. 1908. 40c.
- 131 — Report of the State Botanist 1908. 202p. 4pl. July 1909. 40c.
- 139 — Report of the State Botanist 1909. 116p. 10pl. May 1910. 45c.
- Archeology. 16 Beauchamp, W. M. Aboriginal Chipped Stone Implements of New York. 86p. 23pl. Oct. 1897. 25c.
- 18 — Polished Stone Articles used by the New York Aborigines. 104p. 35pl. Nov. 1897. 25c.
- 22 — Earthenware of the New York Aborigines. 78p. 33pl. Oct. 1898. 25c.
- 32 — Aboriginal Occupation of New York. 190p. 16pl. 2 maps Mar. 1900. 30c.
- 41 — Wampum and Shell Articles used by New York Indians. 166p. 28pl. Mar. 1901. 30c.
- 50 — Horn and Bone Implements of the New York Indians. 112p. 43pl. Mar. 1902. 30c.
- 55 — Metallic Implements of the New York Indians. 94p. 38pl. June 1902. 25c.
- 73 — Metallic Ornaments of the New York Indians. 122p. 37pl. Dec. 1903. 30c.
- 78 — History of the New York Iroquois. 340p. 17pl. map. Feb. 1905. 75c. cloth.
- 87 — Perch Lake Mounds. 84p. 12pl. Apr. 1905. 20c.
- 89 — Aboriginal Use of Wood in New York. 190p. 35pl. June 1905. 35c.
- 108 — Aboriginal Place Names of New York. 336p. May 1907. 40c.
- 113 — Civil, Religious and Mourning Councils and Ceremonies of Adoption. 118p. 7pl. June 1907. 25c.
- 117 Parker, A. C. An Erie Indian Village and Burial Site. 102p. 38pl. Dec. 1907. 30c.
- 125 Converse, H. M. & Parker, A. C. Iroquois Myths and Legends. 196p. il. 11pl. Dec. 1908. 50c.
- Miscellaneous. Ms 1 (62) Merrill, F. J. H. Directory of Natural History Museums in United States and Canada. 236p. Apr. 1903. 30c.
- 66 Ellis, Mary. Index to Publications of the New York State Natural History Survey and New York State Museum 1837-1902. 418p. June 1903. 75c. cloth.
- Museum memoirs 1889-date. Q.
- 1 Beecher, C. E. & Clarke, J. M. Development of Some Silurian Brachiopoda. 96p. 8pl. Oct. 1889. \$1.
- 2 Hall, James & Clarke J. M. Paleozoic Reticulate Sponges. 350p. il. 70pl. 1898. \$2. cloth.
- 3 Clarke, J. M. The Oriskany Fauna of Becraft Mountain, Columbia Co., N. Y. 128p. 9pl. Oct. 1900. 80c.
- 4 Peck, C. H. N. Y. Edible Fungi, 1895-99. 106p. 25pl. Nov. 1900. [\$1.25]
- This includes revised descriptions and illustrations of fungi reported in the 49th, 51st and 52d reports of the State Botanist.

NEW YORK STATE EDUCATION DEPARTMENT

- 5 Clarke, J. M. & Ruedemann, Rudolf. Guelph Formation and Fauna of New York State. 196p. 21pl. July 1903. \$1.50, *cloth*.
- 6 Clarke, J. M. Naples Fauna in Western New York. 268p. 26pl. map. \$2, *cloth*.
- 7 Ruedemann, Rudolf. Graptolites of New York. Pt 1 Graptolites of the Lower Beds. 350p. 17pl. Feb. 1905. \$1.50, *cloth*.
- 8 Felt, E. P. Insects Affecting Park and Woodland Trees. v.1. 460p. il. 48pl. Feb. 1906. \$2.50, *cloth*; v.2. 548p. il. 22pl. Feb. 1907. \$2, *cloth*.
- 9 Clarke, J. M. Early Devonian of New York and Eastern North America. Pt 1. 366p. il. 70pl. 5 maps. Mar. 1908. \$2.50, *cloth*; Pt 2, 250p. il. 36pl. 4 maps. Sept. 1909. \$2, *cloth*.
- 10 Eastman, C. R. The Devonian Fishes of the New York Formations. 236p. 15pl. 1907. \$1.25, *cloth*.
- 11 Ruedemann, Rudolf. Graptolites of New York. Pt 2 Graptolites of the Higher Beds. 584p. il. 2 tab. 31pl. Apr. 1908. \$2.50, *cloth*.
- 12 Eaton, E. H. Birds of New York. v. 1, 501p. il. 42 pl. Apr. 1910. \$3, *cloth*; v. 2 *in press*.
- 13 Whitlock, H. P. Calcites of New York. *In press*.
Clarke, J. M. & Ruedemann, Rudolf. The Eurypterida of New York. *Prepared*.

Natural history of New York. 30v. il. pl. maps. 4to. Albany 1842-04.
DIVISION 1 ZOOLOGY. De Kay, James E. Zoology of New York; or, The New York Fauna; comprising detailed descriptions of all the animals hitherto observed within the State of New York with brief notices of those occasionally found near its borders, and accompanied by appropriate illustrations. 5v. il. pl. maps. sq. 4to. Albany 1842-44. *Out of print*.
Historical introduction to the series by Gov. W. H. Seward. 178p.

- v. 1 pt1 Mammalia. 131+46p. 33pl. 1842.
300 copies with hand-colored plates.
- v. 2 pt2 Birds. 12+380p. 141pl. 1844.
Colored plates.
- v. 3 pt3 Reptiles and Amphibia. 7+98p. pt4 Fishes. 15+415p. 1842.
pt3-4 bound together.
- v. 4 Plates to accompany v. 3. Reptiles and Amphibia 23pl. Fishes 79pl. 1842.
300 copies with hand-colored plates.
- v. 5 pt5 Mollusca. 4+271p. 40pl. pt6 Crustacea. 70p. 13pl. 1843-44.
Hand-colored plates; pt5-6 bound together.

DIVISION 2 BOTANY. Torrey, John. Flora of the State of New York; comprising full descriptions of all the indigenous and naturalized plants hitherto discovered in the State, with remarks on their economical and medical properties. 2v. il. pl. sq. 4to. Albany 1843. *Out of print*.

- v. 1 Flora of the State of New York. 12+484p. 72pl. 1843.
300 copies with hand-colored plates.
- v. 2 Flora of the State of New York. 572p. 89pl. 1843.
300 copies with hand-colored plates.

DIVISION 3 MINERALOGY. Beck, Lewis C. Mineralogy of New York; comprising detailed descriptions of the minerals hitherto found in the State of New York, and notices of their uses in the arts and agriculture. il. pl. sq. 4to. Albany 1842. *Out of print*.

- v. 1 pt1 Economical Mineralogy. pt2 Descriptive Mineralogy. 24+536p. 1842.
8 plates additional to those printed as part of the text.

DIVISION 4 GEOLOGY. Mather, W. W.; Emmons, Ebenezer; Vanuxem, Lardner & Hall, James. Geology of New York. 4v. il. pl. sq. 4to. Albany 1842-43. *Out of print*.

- v. 1 pt1 Mather, W. W. First Geological District. 37+653p. 46pl. 1843.
- v. 2 pt2 Emmons, Ebenezer. Second Geological District. 10+437p. 17pl. 1842.
- v. 3 pt3 Vanuxem, Lardner. Third Geological District. 306p. 1842.
- v. 4 pt4 Hall, James. Fourth Geological District. 22+683p. 19pl. map. 1843.

MUSEUM PUBLICATIONS

DIVISION 5 AGRICULTURE. Emmons, Ebenezer. Agriculture of New York; comprising an account of the classification, composition and distribution of the soils and rocks and the natural waters of the different geological formations, together with a condensed view of the meteorology and agricultural productions of the State. 5v. il. pl. sq. 4to. Albany 1846-54. *Out of print.*

v. 1 Soils of the State, their Composition and Distribution. 11 + 371p. 21pl. 1846.

v. 2 Analysis of Soils, Plants, Cereals, etc. 8 + 343 + 46p. 42pl. 1849.
With hand-colored plates.

v. 3 Fruits, etc. 8 + 340p. 1851.

v. 4 Plates to accompany v. 3. 95pl. 1851.
Hand-colored.

v. 5 Insects Injurious to Agriculture. 8 + 272p. 50pl. 1854.
With hand-colored plates.

DIVISION 6 PALEONTOLOGY. Hall, James. Paleontology of New York. 8v. il. pl. sq. 4to. Albany 1847-94. *Bound in cloth.*

v. 1 Organic Remains of the Lower Division of the New York System. 23 + 338p. 99pl. 1847. *Out of print.*

v. 2 Organic Remains of Lower Middle Division of the New York System. 8 + 362p. 104pl. 1852. *Out of print.*

v. 3 Organic Remains of the Lower Helderberg Group and the Oriskany Sandstone. pt1, text. 12 + 532p. 1859. [\$3.50]
— pt2. 143pl. 1861. [\$2.50]

v. 4 Fossil Brachiopoda of the Upper Helderberg, Hamilton, Portage and Chemung Groups. 11 + 1 + 428p. 69pl. 1867. \$2.50.

v. 5 pt1 Lamellibranchiata 1. Monomyaria of the Upper Helderberg, Hamilton and Chemung Groups. 18 + 268p. 45pl. 1884. \$2.50.

— — — Lamellibranchiata 2. Dimyaria of the Upper Helderberg, Hamilton, Portage and Chemung Groups. 62 + 293p. 51pl. 1885. \$2.50.

— pt2 Gasteropoda, Pteropoda and Cephalopoda of the Upper Helderberg, Hamilton, Portage and Chemung Groups. 2v. 1879. v. 1, text. 15 + 402p. v. 2, 120pl. \$2.50 for 2 v.

— & Simpson, George B. v. 6 Corals and Bryozoa of the Lower and Upper Helderberg and Hamilton Groups. 24 + 208p. 67pl. 1887. \$2.50.

— & Clarke, John M. v. 7 Trilobites and other Crustacea of the Oriskany, Upper Helderberg, Hamilton, Portage, Chemung and Catskill Groups. 64 + 236p. 46pl. 1888. Cont. supplement to v. 5, pt2. Pteropoda, Cephalopoda and Annelida. 42p. 18pl. 1888. \$2.50.

— & Clarke, John M. v. 8 pt1 Introduction to the Study of the Genera of the Paleozoic Brachiopoda. 16 + 367p. 44pl. 1892. \$2.50.

— & Clarke, John M. v. 8 pt2 Paleozoic Brachiopoda. 16 + 394p. 64pl. 1894. \$2.50.

Catalogue of the Cabinet of Natural History of the State of New York and of the Historical and Antiquarian Collection annexed thereto. 242p. 8vo. 1853.

Handbooks 1893-date.

In quantities. 1 cent for each 16 pages or less. Single copies postpaid as below.

New York State Museum. 52p. il. Free.

Outlines, history and work of the museum with list of staff 1902.

Paleontology. 12p. Free.

Brief outline of State Museum work in paleontology under heads: Definition Relation to biology; Relation to stratigraphy; History of paleontology in New York.

Guide to Excursions in the Fossiliferous Rocks of New York. 124p. Free.

Itineraries of 37 trips covering nearly the entire series of Paleozoic rocks, prepared specially for the use of teachers and students desiring to acquaint themselves more intimately with the classic rocks of this State

Entomology. 16p. Free.


Economic Geology. 44p. Free.

Insecticides and Fungicides. 20p. Free

Classification of New York Series of Geologic Formations. 32p. Free.

NEW YORK STATE EDUCATION DEPARTMENT

- Geologic maps.** Merrill, F. J. H. Economic and Geologic Map of the State of New York; issued as part of Museum bulletin 15 and 48th Museum Report, v. 1. 59x67 cm. 1894. Scale 14 miles to 1 inch. 15c.
- Map of the State of New York Showing the Location of Quarries of Stone Used for Building and Road Metal. Mus. bul. 17. 1897. Free.
- Map of the State of New York Showing the Distribution of the Rocks Most Useful for Road Metal. Mus. bul. 17. 1897. Free.
- Geologic Map of New York. 1901. Scale 5 miles to 1 inch. *In atlas form* \$3; *mounted on rollers* \$5. *Lower Hudson sheet* 60c.
- The lower Hudson sheet, geologically colored, comprises Rockland, Orange, Dutchess, Putnam, Westchester, New York, Richmond, Kings, Queens and Nassau counties, and parts of Sullivan, Ulster and Suffolk counties; also northeastern New Jersey and part of western Connecticut.
- Map of New York Showing the Surface Configuration and Water Sheds. 1901. Scale 12 miles to 1 inch. 15c.
- Map of the State of New York Showing the Location of its Economic Deposits. 1904. Scale 12 miles to 1 inch. 15c.
- Geologic maps on the United States Geological Survey topographic base; scale 1 in. = 1 m. Those marked with an asterisk have also been published separately.
- *Albany county. Mus. rep't 49, v. 2. 1898. *Out of print.*
- Area around Lake Placid. Mus. bul. 21. 1898.
- Vicinity of Frankfort Hill [parts of Herkimer and Oneida counties]. Mus. rep't 51, v. 1. 1899.
- Rockland county. State geol. rep't 18. 1899.
- Amsterdam quadrangle. Mus. bul. 34. 1900.
- *Parts of Albany and Rensselaer counties. Mus. bul. 42. 1901. Free.
- *Niagara river. Mus. bul. 45. 1901. 25c.
- Part of Clinton county. State geol. rep't 19. 1901.
- Oyster Bay and Hempstead quadrangles on Long Island. Mus. bul. 48. 1901.
- Portions of Clinton and Essex counties. Mus. bul. 52. 1902.
- Part of town of Northumberland, Saratoga co. State geol. rep't 21. 1903.
- Union Springs, Cayuga county and vicinity. Mus. bul. 69. 1903.
- *Olean quadrangle. Mus. bul. 69. 1903. Free.
- *Becraft Mt with 2 sheets of sections. (Scale 1 in. = $\frac{1}{2}$ m.) Mus. bul. 69. 1903. 20c.
- *Canandaigua-Naples quadrangles. Mus. bul. 63. 1904. 20c.
- *Little Falls quadrangle. Mus. bul. 77. 1905. Free.
- *Watkins-Elmira quadrangles. Mus. bul. 81. 1905. 20c.
- *Tully quadrangle. Mus. bul. 82. 1905. Free.
- *Salamanca quadrangle. Mus. bul. 80. 1905. Free.
- *Mooers quadrangle. Mus. bul. 83. 1905. Free.
- *Buffalo quadrangle. Mus. bul. 99. 1906. Free.
- *Penn Yan-Hammondsport quadrangles. Mus. bul. 101. 1906. 20c.
- *Rochester and Ontario Beach quadrangles. Mus. bul. 114. 20c.
- *Long Lake quadrangle. Mus. bul. 115. Free.
- *Nunda-Portage quadrangles. Mus. bul. 118. 20c.
- *Remsen quadrangle. Mus. bul. 126. 1908. Free.
- *Geneva-Ovid quadrangles. Mus. bul. 128. 1909. 20c.
- *Port Leyden quadrangle. Mus. bul. 135. 1910. Free.
- *Auburn-Genoa quadrangles. Mus. bul. 137. 1910. 20c.
- *Elizabethtown and Port Henry quadrangles. Mus. bul. 138. 1910. 15c.







SMITHSONIAN INSTITUTION LIBRARIES



3 9088 01300 7786